

# MAINE STATE LEGISLATURE

The following document is provided by the  
**LAW AND LEGISLATIVE DIGITAL LIBRARY**  
at the Maine State Law and Legislative Reference Library  
<http://legislature.maine.gov/lawlib>



Reproduced from scanned originals with text recognition applied  
(searchable text may contain some errors and/or omissions)

---

---

# *Annual Report on Air Quality 1987*



## MAINE

Department of  
Environmental Protection



TD  
883.5  
.M2  
M33  
1987

---

---

STATE LAW LIBRARY  
AUGUSTA, MAINE



1987 ANNUAL REPORT  
ON AIR QUALITY  
IN THE STATE OF MAINE

Prepared by  
Jeffrey C. Emery  
Environmental Specialist

Maine Department of Environmental Protection  
Bureau of Air Quality Control  
Division of Air Quality Services

Leighton E. Carver, Division Director  
Dennis L. Keschl, Bureau Director  
Dean C. Marriott, Commissioner



## TABLE OF CONTENTS

	<u>PAGE</u>
TABLE OF CONTENTS	i
LIST OF TABLES	iii
LIST OF FIGURES	v
1. INTRODUCTION	1
1.1 Purpose and Overview	1
1.2 Monitoring Sites	6
1.3 Document Organization	17
1.3.1 Explanation of Data Summary Tables	17
1.3.2 Explanation of Historical Comparison Tables	17
1.3.3 Explanation of Trends Tables	17
2. CARBON MONOXIDE (CO)	20
2.1 Description and Sources	20
2.2 Health and Welfare Effects	20
2.3 Standards	21
2.4 Monitoring	21
3. OZONE (O3)	23
3.1 Description and Sources	23
3.2 Health and Welfare Effects	24
3.3 Standards	24
3.4 Monitoring	24
4. NITROGEN DIOXIDE (NO2)	28
4.1 Description and Sources	28
4.2 Health and Welfare Effects	28
4.3 Standards	28
4.4 Monitoring	28
5. SULFUR DIOXIDE (SO2)	30
5.1 Description and Sources	30
5.2 Health and Welfare Effects	30
5.3 Standards	30
5.4 Monitoring	30
6. PARTICULATES (TSP)	35
6.1 Description and Sources	35
6.2 Health and Welfare Effects	35
6.3 Standards	36
6.4 Monitoring	36
7. LEAD (Pb)	45
7.1 Description and Sources	45
7.2 Health and Welfare Effects	45
7.3 Standards	45
7.4 Monitoring	45

TABLE OF CONTENTS (continued)

	<u>PAGE</u>
8. SULFATES (SO <sub>4</sub> ) AND NITRATES (NO <sub>3</sub> )	49
8.1 Description and Sources	49
8.2 Health and Welfare Effects	49
8.3 Standards	49
8.4 Monitoring	49
9. ATMOSPHERIC DEPOSITION	53
9.1 Description and Sources	53
9.2 Health and Welfare Effects	53
9.3 Standards	53
9.4 Monitoring	53
10. HYDROCARBONS (HC)	55
10.1 Description and Sources	55
10.2 Health and Welfare Effects	55
10.3 Standards	55
10.4 Monitoring	55

## LIST OF TABLES

<u>TABLE</u>	<u>TITLE</u>	<u>PAGE</u>
1-1	National Ambient Air Quality Standards	2
1-2	State of Maine Ambient Air Quality Standards	3
1-3	Number of Ambient Air Quality Violations by Regions	4
1-4	1987 Ambient Air Quality Monitoring Site Directory	8
2-1	1987 Carbon Monoxide Data Summary	22
2-2	Carbon Monoxide Historical Comparisons	22
2-3	Carbon Monoxide Trends	22
3-1	1987 Ozone Data Summary	25
3-2	Ozone Historical Comparisons	26
3-3	Ozone Trends	27
4-1	1987 Nitrogen Dioxide Data Summary	29
5-1	1987 Sulfur Dioxide Data Summary	32
5-2	Sulfur Dioxide Historical Comparisons-Maximum 24-Hour Concentrations of Sulfur Dioxide	33
5-3	Sulfur Dioxide Historical Comparisons-Sites with Violations	34
6-1	1987 Total Suspended Particulates Data Summary	38
6-2	Total Suspended Particulates Historical Comparison-Annual Geometric Means	40
6-3	Total Suspended Particulates Historical Comparisons-Sites With Violations	42
6-4	1987 Fine Particulate Data Summary	44
7-1	1987 Lead Data Summary	46
7-2	1987 Lead Data Summary by Quarters	47
7-3	Lead Historical Comparisons	48
8-1	Sulfate Thresholds for Adverse Health Effects	50
8-2	1987 Sulfate Data Summary	51



LIST OF TABLES (continued)

<u>TABLE</u>	<u>TITLE</u>	<u>PAGE</u>
8-3	1987 Nitrate Data Summary	52
9-1	1987 Atmospheric Deposition Data Summary	54

# LIST OF FIGURES

<u>FIGURE</u>	<u>TITLE</u>	<u>PAGE</u>
1-1	Five Year Trend - Total Suspended Particulates	5
1-2	Five Year Trend - Sulfur Dioxide	7
1-3	Five Year Trend - Ozone	7
1-4	State of Maine, Ambient Air Quality Control Regions	18



## 1. INTRODUCTION

### 1.1 Purpose and Overview

The purpose of this report is to present the air quality monitoring data generated by and for the Maine Department of Environmental Protection, Bureau of Air Quality Control, and to provide a historical perspective from which the significance of that data can be interpreted. Air Quality monitoring measures the concentrations of various pollutants in the ambient air. The monitoring is in response to State and Federal requirements to determine whether the air we breathe is attaining and maintaining National and State Ambient Air Quality Standards which are designed to protect the health and welfare of the public. Federal Primary Standards are intended to protect public health. Federal Secondary Standards are intended to protect public welfare. The State Standards are at least as strict as Federal Standards and in some cases are more strict. The reasoning behind establishing more stringent standards is that generally air quality in Maine is significantly cleaner than in other areas and should remain cleaner. The current Federal and State Standards are presented in Tables 1-1 and 1-2. Table 1-3 is a summary indicating all the violations of ambient air quality standards in the State by regions. Later on in this report those violations will be listed by the sites at which they occurred.

A significant portion of the data collected in the State is collected by industry. The Department has required industry to establish monitoring programs primarily when there are air quality problems associated with the industry, or when an industry is planning to build or expand causing a potential increase in air emissions. The State is still collecting monitoring data for long term trends, special studies and for compliance determinations. Ambient air monitoring by both industry and the State will continue in various regions where necessary until such time as standards are being met.

Included in this section are some figures which depict some of the results of air quality monitoring and control in the State. Figures 1-1 through 1-3 display trends or the lack of a trend which have been occurring at several long term key sites around the State.

Figure 1-1 depicts the annual geometric means for total suspended particulates at several long term sites. Only one site still shows a significant upward trend, the Research Building site in Westbrook. The Westbrook trend continues to be the result of increased development, fugitive emissions from the S. D. Warren facility and a sweeping program that hasn't been comprehensive enough. Some reduction occurred in 1986 but the annual geometric mean in 1987 increased to a new high and exceeds the standard by a significant amount. The annual geometric mean at this site has exceeded the state standard for the last four years. The Westbrook area will need increased control efforts to achieve compliance.

TABLE 1-1  
NATIONAL AMBIENT AIR QUALITY STANDARDS  
(1987)

<u>Pollutant</u>	<u>Averaging Time</u>	<u>Concentration</u>
Particulates (TSP)	Annual Geometric Mean:	
	Primary	75 ug/m3
	Secondary	60 ug/m3 *
	Twenty-Four Hour:**	
	Primary	260 ug/m3
	Secondary	150 ug/m3
Particulates (PM10)	Expected Annual Arithmetic Mean:	
	Primary	50 ug/m3
	Secondary	50 ug/m3
	Twenty-Four Hour:***	
	Primary	150 ug/m3
	Secondary	150 ug/m3
Lead (Pb)	Calendar Quarter	1.5 ug/m3
Carbon Monoxide (CO)	One Hour**	35 ppm
	Eight Hour**	9 ppm
Ozone (O3)	One Hour***	0.12 ppm
Nitrogen Dioxide (NO2)	Annual Arithmetic Mean	0.05 ppm
Sulfur Dioxide (SO2)	Annual Arithmetic Mean	0.03 ppm
	Twenty-Four Hour**	0.14 ppm
	Three-Hour**	
	Secondary	0.50 ppm
Hydrocarbon	Three Hour**	160 ug/m3

\* = Federal Guideline Only.

\*\* = Not to be exceeded more than once per year.

\*\*\* = Statistically estimated number of days with exceedances is not to be more than 1 per year.

ppm = Parts of pollutant per million parts of air.

ug/m3 = Micrograms of pollutant per cubic meter of air.

TABLE 1-2  
STATE OF MAINE AMBIENT AIR QUALITY STANDARDS  
(1987)

<u>Pollutant</u>	<u>Averaging Time</u>	<u>Concentration</u>
Particulates (TSP)	Annual Geometric Mean	60 ug/m3
	Twenty-Four Hour	150 ug/m3
Lead (Pb)	Twenty-Four Hour*	1.5 ug/m3
Carbon Monoxide (CO)	One Hour*	35 ppm(40 mg/m3)
	Eight Hour*	9 ppm(10 mg/m3)
Ozone (O3)	One Hour*	.081 ppm(160 ug/m3)
Nitrogen Dioxide (NO2)	Annual Arithmetic Mean	.053 ppm(100 ug/m3)
Sulfur Dioxide (SO2)	Annual Arithmetic Mean	.022 ppm( 57 ug/m3)
	Twenty-Four Hour	.088 ppm(230 ug/m3)
	Three Hour	.439 ppm(1150 ug/m3)
Hydrocarbon	Three Hour*	160 ug/m3

\* = Not to be exceeded more than once per year.  
 PPM = Parts of pollutant per million parts of air.  
 ug/m3 = Micrograms of pollutant per cubic meter of air.  
 mg/m3 = Milligrams of pollutant per cubic meter of air.

TABLE 1-3  
NUMBER OF AMBIENT AIR QUALITY VIOLATIONS BY REGIONS  
(1987)


<u>POLLUTANT</u>	<u>REGIONS</u>					<u>TOTALS</u>
	<u>107</u>	<u>108</u>	<u>109</u>	<u>110</u>	<u>111</u>	
Total Suspended Particulates						
Annual Geometric Mean*						
State	0	0	0	2	?	2
Federal	0	0	0	0	?	0
Twenty-four Hour						
State	8	12	22	19	?	61
Federal	0	2	0	0	?	2
Fine Particulate - Federal						
Annual Arithmetic Mean	0	0	0	0	?	0
Twenty-four Hour	0	2	0	0	?	2
Lead						
Twenty-four Hour						
State	0	0	0	0	?	0
Federal	0	0	0	0	?	0
Carbon Monoxide						
One Hour	?	?	?	0	?	0
Eight Hour	?	?	?	0	?	0
Ozone						
One Hour						
State	194	?	79	197	40	510
Days						
Federal	5	?	0	7	0	12
Nitrogen Dioxide						
Annual Arithmetic Mean	?	?	?	0	?	0
Sulfur Dioxide						
Annual Arithmetic Mean						
State	0	0	0	0	?	0
Federal	0	0	0	0	?	0
Twenty-four Hour						
State	1	0	0	0	?	1
Federal	0	0	0	0	?	0
Three Hour						
State	0	0	0	0	?	0
Federal	0	0	0	0	?	0

\* Annual Means generated by only a few samples are not included in this summary.

? No monitoring done for this pollutant within this region during 1987.

FIGURE 1 - 1

FIVE YEAR TREND - TOTAL SUSPENDED PARTICULATES





Presque Isle has continued to show improvement and is showing a very significant downward trend over the last four years. The overall air quality has improved a lot, but short term violations continue to occur and additional efforts may be needed to bring the area into compliance. The use of a cleaner sand on winter roads, increased clean up efforts during the Spring and a bypass for Presque Isle to keep through traffic out of the downtown area may be the only solution to the particulate problem.

Figure 1-2 indicates the sulfur dioxide trends at three sites with a long term history. All three sites appear to indicate relatively stable sulfur dioxide levels over the last four to five years.

Figure 1-3 depicts the number of hourly violations of the State ozone standard. As can be seen from the graphs, the violations vary greatly from year to year and do show a very significant reduction in the number of violations during 1986 with only a slight increase in 1987. However, meteorological conditions are responsible for a lot of the variability from year to year so it is too early to tell if existing state applied control strategies are having any significant effect on the basic ozone problem.

Data summarized in this report is available for review in the Department headquarters in Augusta and copies can be obtained from that office for a nominal fee.

## 1.2 Monitoring Sites

Air quality data are developed using two basic methods; 1) the continuous monitoring of gaseous pollutants and; 2) the periodic sampling of particulate and gaseous pollutants. In addition to pollutant monitoring there is also the continuous monitoring of meteorological parameters.

Continuous gaseous monitoring was done at thirty-four sites in Maine during 1987. Carbon Monoxide was monitored at one of these stations, ozone at ten, nitrogen dioxide at two and sulfur dioxide at eighteen.

Particulate sampling was done at fifty-four sites in Maine during 1987. Fifty of these stations monitored total suspended particulates. Sixteen of these sites also collected fine particulate fractions. Also, lead monitoring was done at seven stations. Six sites were analyzed for sulfates and nitrates. There were also two sites collecting acid rain data as part of the state monitoring network.

In addition to pollutant monitoring, wind speed and direction was recorded at twenty-three sites around the State during 1987. Some of these sites also recorded other meteorological parameters such as sigma (stability) and temperature, precipitation and solar radiation.

Table 1-4 presents all the monitoring sites in Maine that operated during 1987 and indicates which parameters were monitored at

FIGURE 1 - 2  
FIVE YEAR TREND - SULFUR DIOXIDE




FIGURE 1 - 3  
FIVE YEAR TREND - OZONE




TABLE 1 - 4  
1987 AMBIENT AIR QUALITY MONITORING SITE DIRECTORY

<u>SITE</u>	<u>ADDRESS</u>	<u>OPERATOR</u>	<u>PARAMETERS MEASURED</u>
<b>ANDROSCOGGIN INTERSTATE AIR QUALITY CONTROL REGION (107)</b>			
Auburn (0060 005)	Lewiston-Auburn Airport Lewiston Junction Road	DEP	WS/WD
Auburn (0060 008)	Lepage Bakery 60 Second Street	DEP	TSP,Pb
Augusta (0080 005)	Hartford Fire House Hartford Square	DEP	TSP,FP
Augusta (0080 008)	Governor's Hangar State Airport	DEP	WS/WD
Gardiner (0460 001)	Gardiner High School West Hill Road	DEP	Ozone(s)
Jay (0530 001)	Weather Level I Lagoon Hill	International Paper	WS/WD, Temperature, Solar Radiation, Precipitation, TSP
Jay (0530 003)	Crash Road Gilbert Jewell Property	International Paper	TSP
Jay (0530 004)	Jay Hill	International Paper	TSP,FP
Jay (0530 008)	Burnham Site	International Paper	TSP
Port Clyde(NEW) (0595 004)	Port Clyde Ozone St. George	DEP	Ozone(s)
Isle Au Haut (0595 003)	Isle Au Haut Fire Station	UM/DEP	Ozone(s)

TABLE 1 - 4 (continued)  
1987 AMBIENT AIR QUALITY MONITORING SITE DIRECTORY

<u>SITE</u>	<u>ADDRESS</u>	<u>OPERATOR</u>	<u>PARAMETERS MEASURED</u>
Lewiston (0620 011)	Country Kitchen Parking Lot Canal Street	DEP	SO2
Mexico (0760 008)	Labonville's Route #2	Boise Cascade	TSP,FP(n)
Mexico (0760 011)	Hunt's Property Route #2	Boise Cascade	SO2
Rumford (1020 002)	Boise Cascade Weather II Swift River Pump House	Boise Cascade	WS/WD
Rumford (1020 005)	Taylor Mountain I	Boise Cascade	TSP,SO2,Sulfate,Nitrate
Rumford (1020 006)	Taylor Mountain II	Boise Cascade	TSP,SO2
Rumford (1020 007)	Village Green Site Route #108	DEP/Boise Cascade	TSP,SO2
Skowhegan (1100 001)	Hinckley Hinckley Farm School	S. D. Warren	TSP
Skowhegan (1100 002)	Eaton Ridge	S. D. Warren	TSP
Thomaston (1150 001)	Mitchell Property 2 Dexter Avenue	Dragon Products	TSP
Thomaston (1150 003)	Sanders Property Old County Road	Dragon Products	TSP
Thomaston (1150 004)	Pease Heirs Property Buttermilk Lane	Dragon Products	TSP

TABLE 1 - 4 (continued)  
1987 AMBIENT AIR QUALITY MONITORING SITE DIRECTORY

<u>SITE</u>	<u>ADDRESS</u>	<u>OPERATOR</u>	<u>PARAMETERS MEASURED</u>
Thomaston (1150 005)	Dragon Cement Weather Route #1	Dragon Products	WS/WD
Thomaston (1150 007)	Marsh Road	Dragon Products	TSP
Searsport (1183 006)	DOT Route #1	DEP	SO2,WS/WD
Waterville (1220 003)	Stern's Department Store Main Street	DEP	TSP
Winslow (1280 003)	Gulley Hill Road	Scott Paper Company	TSP,FP

**AROOSTOOK AIR QUALITY CONTROL REGION (108)**

Fort Fairfield(DISC) (0400 001)	Peterson's	Fairfield Energy Venture	TSP
Fort Kent(NEW) (0420 001)	UMFK - Cyr Hall	UMFK	TSP,WS/WD
Madawaska (0720 003)	Madawaska High School 7th Avenue	Fraser Paper	SO2
Madawaska (0720 006)	Fraser Paper Company Bridge Street	Fraser Paper	WS/WD, Temperature
Madawaska (0720 009)	Albert Street	Fraser Paper	SO2
Madawaska (0720 011)	St. Jarre's 11th Avenue	DEP	TSP, Sulfate, Nitrate

TABLE 1 - 4 (continued)  
1987 AMBIENT AIR QUALITY MONITORING SITE DIRECTORY

<u>SITE</u>	<u>ADDRESS</u>	<u>OPERATOR</u>	<u>PARAMETERS MEASURED</u>
Madawaska (0720 012)	U. S. Post Office 430 E. Main Street	Fraser Paper	SO <sub>2</sub> ,WS/WD
Madawaska (0720 013)	Big Daddy's Restaurant 395 E. Main Street	DEP	FP
Presque Isle (0980 005)	Northeastland Hotel 436 Main Street	DEP	TSP,Pb,FP
Presque Isle (0980 008)	Regional Office 528 Central Drive	DEP	WS/WD

**DOWNEAST AIR QUALITY CONTROL REGION (109)**

Acadia National Park (0010 003)	McFarland Hill Ranger Station Route #233	NPS/DEP	Ozone,TSP,Sulfate,Nitrate,FP,Acid Precipitation
Bangor (0100 001)	Regional Office 31 Central Street	DEP	TSP,Sulfate,Nitrate
Bangor (0100 002)	Kenduskeag Pump Station Washington Street	DEP	TSP,Pb,FP,SO <sub>2</sub> (d)
Bangor(DISC) (0100 009)	BIA-Building #487 Air National Guard	DEP	WS/WD
Bangor(NEW) (0100 010)	BIA-Building #489 Air National Guard	DEP	WS/WD
Bangor(NEW) (0100 011)	Flight Service Station Bangor International Airport	FAA	Temperature
Brewer (0180 002)	Brewer Junior High School 5 Somerset Street	DEP	TSP

TABLE 1 - 4 (continued)  
1987 AMBIENT AIR QUALITY MONITORING SITE DIRECTORY

<u>SITE</u>	<u>ADDRESS</u>	<u>OPERATOR</u>	<u>PARAMETERS MEASURED</u>
Bucksport (0205 003)	Bucks Mill Road	Champion International	TSP(d),WS/WD
Bucksport(DISC) (0205 004)	Public Landing	Champion International	TSP,WS/WD
Bucksport(NEW) (0205 005)	Waste Disposal Site Route #15	Champion International	WS/WD, Temperature, Precipitation
Calais (0220 003)	WOODY Tower	Georgia Pacific Corporation	WS/WD
Hampden(NEW) (0485 001)	McGraw School	Penobscot Energy Recovery Company	FP,Pb,Chrome,Heavy Metals
East Millinocket (0315 002)	Katahdin School School Street	Great Northern Paper Company	TSP,S02(i)
Dedham (0495 003)	Bald Mountain	DEP	Ozone(s),WS/WD(s)
Lincoln (0640 002)	Vocational Education Building West Broadway	Lincoln Pulp & Paper Company	TSP
Lincoln (0640 003)	Lincoln Post Office Building 50 Fleming Street	Lincoln Pulp & Paper Company	TSP
Lincoln (0640 007)	Thomas Motel Trailer Park 39 West Broadway	Lincoln Pulp & Paper Company	TSP,S02,FP
Lincoln (0640 008)	Fish Hill Base	Lincoln Pulp & Paper Company	S02
Lincoln (0640 009)	Fish Hill Peak	Lincoln Pulp & Paper Company	S02

TABLE 1 - 4 (continued)  
1987 AMBIENT AIR QUALITY MONITORING SITE DIRECTORY

<u>SITE</u>	<u>ADDRESS</u>	<u>OPERATOR</u>	<u>PARAMETERS MEASURED</u>
Lincoln (0640 010)	Lincoln Airport	Lincoln Pulp & Paper Company	WS/WD
Millinocket (0780 006)	Wastewater Treatment Plant Great Northern Paper Company	Great Northern Paper Company	SO2
Millinocket (0780 009)	York Street	Great Northern Paper Company	TSP,SO2
Millinocket (0780 011)	Great Northern Paper Co. Office	Great Northern Paper Company	WS/WD
Old Town (0840 003)	Marsh Island Apartments 100 South Main Street	DEP	TSP
Old Town(DISC) (0840 005)	Penobscot Shoe Company 450 North Main Street	DEP	TSP
Orrington(NEW) (0845 005)	Center Drive School	Penobscot Energy Recovery Company	FP,Pb,Chrome,Heavy Metals
Newburgh (0907 005)	Newburgh School Route #9	DEP	TSP
Milford (0907 007)	Shumway Field Route #178	James River Corporation	TSP
Woodland (1205 007)	Secondary Treatment Pipeline	Georgia Pacific Corporation	TSP,SO2
Woodland (1205 008)	Woodland High School	Georgia Pacific Corporation	TSP,FP
Woodland (1205 017)	Woodyard Woodland Mill	Georgia Pacific Corporation	WS/WD,Temperature



TABLE 1 - 4 (continued)  
1987 AMBIENT AIR QUALITY MONITORING SITE DIRECTORY

<u>SITE</u>	<u>ADDRESS</u>	<u>OPERATOR</u>	<u>PARAMETERS MEASURED</u>
Woodland (1205 018)	Background	Georgia Pacific Corporation	TSP
<b>METROPOLITAN PORTLAND AIR QUALITY CONTROL REGION (110)</b>			
Berwick (0150 001)	Berwick Fire Station Berwick	DEP	TSP
Biddeford (0160 002)	Biddeford Treatment Plant Water Street	DEP	TSP
Bridgton (0190 002)	Upper Ridge Road	DEP	Acid Precipitation,TSP,Sulfate,Nitrate,FP
Brunswick(NEW) (0200 003)	Cook's Corner	DEP	TSP
Cape Elizabeth (0250 003)	Shelter Site Two Lights State Park	DEP	Ozone(s),WS/WD,NO(n),NOX(n)
Portland (0960 010)	Chevrus High School Ocean Avenue	DEP	WS/WD
Portland (0960 014)	Shelter Site (P.E.O.P.L.) Elm Street	DEP	TSP,Pb,SO2,FP,NO(n),NOX(n)
Portland (0960 018)	Congress Street	DEP	CO
Portland(NEW) (0960 019)	YWCA 87 Spring Street	DEP	TSP,Pb
Portland(NEW) (0960 020)	Elks Lodge 1945 Congress Street	Regional Waste Systems	FP

TABLE 1 - 4 (continued)  
1987 AMBIENT AIR QUALITY MONITORING SITE DIRECTORY

<u>SITE</u>	<u>ADDRESS</u>	<u>OPERATOR</u>	<u>PARAMETERS MEASURED</u>
South Portland (1140 002)	SMVTI Vocational Drive	DEP	TSP,Sulfate,Nitrate
Westbrook (1260 002)	N. E. T. & T. Company Ash Street	S. D. Warren	TSP
Westbrook (1260 008)	Research Building S. D. Warren	S. D. Warren	TSP
Westbrook (1260 009)	S. D. Warren Company Wind S. D. Warren Property	S. D. Warren	WS/WD
Westbrook (1260 012)	S. D. Warren Warehouse #5 Main Street	S. D. Warren	TSP,FP
Cape Neddick (1315 001)	Cape Neddick Light Station	DEP	Ozone(s)
Kennebunkport (1325 002)	Parson's Way	DEP	Ozone(s)

**NORTHWEST MAINE AIR QUALITY CONTROL REGION (111)**

T4R2 BKP WKR (0445 001)	Sugarloaf Mountain	UM/DEP	Ozone(s)
T4R2 BKP WKR (0445 002)	Sugarloaf Mountain - Base	DEP	Ozone(s)
Greenville (0935 001)	Squaw Brook Greenville	DEP	Acid Precipitation

TABLE 1 - 4 (continued)  
1987 AMBIENT AIR QUALITY MONITORING SITE DIRECTORY

<u>SITE</u>	<u>ADDRESS</u>	<u>OPERATOR</u>	<u>PARAMETERS MEASURED</u>
	NEW - Site established in 1987		n - Instrument installed during 1987
	DISC - Site discontinued in 1987		d - Instrument removed during 1987
			s - Instrument operated seasonally during 1987
			i - Instrument operated intermittently during 1987
	TSP - Total Suspended Particulates		
	SO2 - Sulfur Dioxide		
	NO - Nitric Oxide		
	NOX - Oxides of Nitrogen		
	CO - Carbon Monoxide		
	Pb - Lead		
	WS/WD - Wind Speed and Direction		
	FP - Fine Particulate		
	NMHC - Nonmethane Hydrocarbons		

each site. The map in Figure 1-4 shows the Air Quality Control Regions within the State.

### **1.3 Document Organization**

This document is divided by pollutant into chapters. Each chapter contains: 1) a description of the nature and sources of that pollutant, 2) its health and welfare effects, 3) a discussion on the standards (current and proposed) for that pollutant, 4) a discussion of the monitoring methods for that pollutant, 5) a table presenting the 1987 monitored data, 6) in the case of some pollutants, historical tables presenting 1987 data along with data for previous years to show trends, effects of control strategy, or change in emission sources.

#### **1.3.1 Explanation of Data Summary Tables**

The Data Summary Tables were designed to facilitate comparing 1987 air quality monitoring data with the standards for each pollutant. Therefore, the data are presented for each averaging time for which standards exist for a pollutant.

An annual average concentration is presented for each pollutant that has a long-term, annual standard (NO<sub>2</sub>, SO<sub>2</sub>, TSP).

For pollutants that have short-term standards, the highest short-term values are presented. Some pollutants are allowed to exceed the standard once during the year so the second highest value would be used to determine whether there was a violation or not.

All of the data collected during 1987 has been presented in the Data Summary Tables. However, in making comparisons of the data, one should be aware that a site with only a few samples will not be a valid indicator of pollutant concentrations in the area.

#### **1.3.2 Explanation of Historical Comparison Tables**

The Historical Comparison Tables present air quality data for 1987 and those years prior to 1987 when the same pollutant was monitored at the same site. The purpose of the Tables are to indicate the variations in air quality from year to year. The Tables in some cases represent maximum concentrations for specific time periods and in others the number of days in each year that the standards were violated.

#### **1.3.3 Explanation of Trends Tables**

The highest hourly concentration in a year is not the best indicator of long-term air quality trends because it is an erratic value. Therefore, special trend tables are presented for carbon monoxide and ozone. The trend tables present the 10th, 50th, and 90th percentile values to represent the bulk of the air quality data for each year. Percentiles indicate the fraction, or percent, of the value that are below a particular level. For example, if the 90th

Northwest Maine  
Air Quality Control  
Region(111)

- Arrostook Air  
Quality Control  
Region(108)

Downeast Air  
Quality Control  
Region(109)

Androscoggin Interstate  
Air Quality Control Region(107)

Metropolitan Portland Air  
Quality Control Region(110)

FIGURE 1-4: AIR QUALITY CONTROL REGIONS  
IN THE STATE OF MAINE

MINOR CIVIL DIVISIONS

STATE OF  
**MAINE**

PREPARED BY THE  
STATE OF MAINE  
DEPARTMENT OF TRANSPORTATION  
BUREAU OF PLANNING  
IN COOPERATION WITH THE  
U.S. DEPARTMENT OF TRANSPORTATION  
FEDERAL HIGHWAY ADMINISTRATION

1980

percentile value for some set of CO observations is 5.0 ppm, it means that 90% of the time the concentrations of CO are less than 5.0 ppm. Conversely, it also means that 10% of the time the concentrations are above 5.0 ppm. Thus the existence or lack of long-term trends in overall air quality for CO and O3 can be more reliably determined using the Trends Tables, than by looking at just the Historical Comparison Tables.

## 2. CARBON MONOXIDE (CO)

### 2.1 Description and Sources

Carbon monoxide is colorless, odorless and tasteless gas. Therefore you do not even know you are breathing it until you feel its detrimental effects. It constitutes the largest single fraction of the pollutants found in urban atmospheres. It is produced primarily by the incomplete combustion of organic materials used as fuels for transportation and in the heating of buildings; it also results from industrial processes, refuse burning, and agricultural burning. Several natural sources of CO of both biological and non-biological origin have also been identified, but their contributions to urban atmospheric concentrations are thought to be small. Background levels of CO (resulting from natural and technological sources) found in relatively nonpolluted air range from 0.025 to 1.0 ppm. Urban carbon monoxide is produced primarily by motor vehicles.

Because motor vehicle traffic is the major source of CO, daily concentration peaks coincide with morning and evening rush hours. The worst carbon monoxide problems are found where large numbers of slow moving cars congregate. These problems are further aggravated when they occur in a "street canyon" situation. When there are large amounts of slow moving traffic in a street canyon situation, with the wind blowing perpendicular to the street, carbon monoxide can be trapped in the canyon and build up to unhealthful levels.

CO problems are usually worse in winter because: 1) cold weather makes motor vehicles run dirtier and requires more combustion for space heating; and 2) on winter nights a strong inversion layer develops in the atmosphere, that traps pollution near the ground, preventing it from mixing with cleaner air above.

### 2.2 Health and Welfare Effects

Carbon monoxide affects the central nervous system by depriving the body of the oxygen it needs. Tests of automobile drivers show exposure to carbon monoxide can impair driver's judgement and ability to respond rapidly in traffic. It can also impair vision and produce headaches.

Carbon monoxide enters the bloodstream by combining with hemoglobin, the substance that carries oxygen to the cells. Hemoglobin that is bound up with CO is called carboxyhemoglobin. This combination occurs 200 times more readily with CO than with oxygen, so the amount of oxygen being distributed throughout the body by the bloodstream is reduced in CO's presence. Blood laden with CO can weaken heart contractions, lowering the volume of blood distributed to various parts of the body. It can also significantly reduce a healthy person's ability to perform manual tasks, such as working, jogging and walking. A life-threatening situation exists in patients with heart disease, who can't compensate for the oxygen loss. The 4.2

million people in the U.S. suffering from angina pectoris (a heart disease characterized by brief spasmodic attacks of chest pain due to insufficient oxygen levels in the heart muscles) are especially susceptible. Carbon monoxide is also harmful to persons who have lung disease, anemia or cerebral-vascular disease. Others sensitive to carbon monoxide include the human fetus, and people exposed to long-term concentrations, such as traffic officers.

People who sit in idling cars over sustained periods risk harmful CO exposure, as do cigarette smokers. Since about two percent of cigarette smoke is carbon monoxide, if you or someone else smokes while driving in heavy traffic, you may both experience the harmful effects of CO from the cigarette smoke and the engine exhaust accumulated in streets. Even three or four hours after you're exposed, half the excess CO still remains in your bloodstream. Because it takes time for CO to build up in the bloodstream, the severity of health effects depends both on the concentration being breathed and the length of time the person is exposed.

### 2.3 Standards

The existing standards for carbon monoxide are currently set at 9 parts CO per million parts air (ppm), averaged over a period of 8 hours, and 35 ppm averaged over 1 hour, not to be exceeded more than once per year. As a result of a review and revision of the health criteria, EPA proposes to retain the existing primary 8-hour standard at 9 ppm and to lower the primary 1-hour standard to 25 ppm. The change in the 1-hour standard is being proposed because of the more rapid accumulation of blood carboxyhemoglobin in moderately exercising sensitive persons compared to resting individuals. The impact of exercise, which is greater for short-duration exposures, was not considered in the original standard.

### 2.4 Monitoring

Carbon monoxide was monitored at one site in Maine during 1987 using continuous monitoring equipment utilizing the non-dispersive infrared technique.

Table 2-1 is the 1987 Data Summary for CO. Tables 2-2 and 2-3 have been included for historical comparisons and trend analysis.



TABLE 2 - 1  
1987 CARBON MONOXIDE DATA SUMMARY  
(Parts Per Million)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>1-HOUR CONCENTRATIONS HIGHEST</u>	<u>SECOND HIGHEST</u>	<u>8-HOUR CONCENTRATIONS HIGHEST</u>	<u>SECOND HIGHEST</u>	<u>ANNUAL ARITH. MEAN</u>
<b>METROPOLITAN PORTLAND AIR QUALITY CONTROL REGION (110)</b>							
Portland	Congress Street	7235	8.3	7.3	5.7	5.7	1.4

TABLE 2 - 2  
CARBON MONOXIDE HISTORICAL COMPARISONS

PORTLAND Portland-Congress Street		
<u>YEAR</u>	<u>SECOND HIGH*</u>	<u>NUMBER OF VIOLATIONS</u>
1984	6.9	0
1985	5.9	0
1986	5.1	0
1987	5.7	0

\* Eight hour concentrations in ppm.

TABLE 2 - 3  
CARBON MONOXIDE TRENDS

PORTLAND Portland-Congress Street			
<u>YEAR</u>	<u>Percentiles*</u>		
	<u>10%</u>	<u>50%</u>	<u>90%</u>
1984	0.2	1.0	2.9
1985	0.4	1.1	2.9
1986	0.3	1.1	2.7
1987	0.4	1.1	2.6

\* Percentiles are one hour concentrations in ppm.

### 3. OZONE (O3)

#### 3.1 Description and Sources

Ozone is a highly reactive form of oxygen which, at very high concentrations, is a blue unstable gas that has a characteristic pungent odor most commonly identified around an arcing electric motor, lightning storms, or other electrical discharges. However, at normal ambient concentrations, ozone is colorless and odorless. Ozone is the major component of photochemical "smog", but the haziness and odors of smog are primarily caused by other components.

Natural ground level ozone occurs in low concentrations (less than .05 ppm) due to natural physical and chemical phenomena. Occasionally, unique meteorological conditions can result in natural levels between .05 and .10 ppm.

Ozone is not emitted directly from a source as are other pollutants. It forms as secondary pollutant. Its precursors are hydrocarbons and nitrogen oxides, which chemically react in sunlight to form ozone. The hydrocarbons are emitted in automobile exhaust, from gasoline and oil storage and transfer, and from industrial use of paint solvents, degreasing agents, cleaning fluids, ink solvents, incompletely burned coal or wood and many other sources. Plants also give off hydrocarbons such as terpenes from pine trees. Nitrogen oxides are emitted by all combustion sources.

The highest ozone levels generally occur during summer afternoons when the high temperatures and strong sunlight promote photochemical reactions. Stagnant weather may cause smog to remain in an area for several days. The winds may also transport ozone many miles outside of the urban environment. For example, it is estimated that the majority of the ozone in the State of Maine is transported into the State from sources located outside the State. In addition a much smaller amount of the ozone is naturally occurring background concentrations, part of which is also transported into the State. The remaining ozone is assumed to be due to local sources within the State. Because of long-range transport, local control of emissions by itself may not solve the ozone problem. An effective national program may be necessary to achieve national compliance.

Ground-level ozone, discussed above, should not be confused with the stratospheric ozone layer, located about seven miles high in the atmosphere, which shields the earth from cancer-causing ultraviolet rays. Concentrations of ozone in this layer may reach as high as 10 ppm. Concern over potential reduction of the necessary levels of ozone in the stratosphere by reactions with fluorocarbons from aerosol cans has resulted in the removal of most of these propellants from the market. However, ozone at ground level, where it is breathed, is a pollutant.

### 3.2 Health and Welfare Effects

Ozone at low concentrations causes eye irritations and at higher concentrations difficulty in breathing for people with respiratory problems, the elderly, and children. Many plants, such as white pine, soybeans and alfalfa, are extremely sensitive to ozone, and ozone is known to weaken materials such as rubber and fabrics.

### 3.3 Standards

The existing National Ambient Air Quality Standard (NAAQS) for ozone is 0.12 ppm and will be attained when "the expected number of days per calendar year with maximum hourly average concentrations above 0.12 ppm is equal to or less than one". This standard is new as of February 8, 1979 and replaces a more restrictive 0.08 ppm standard that was established April 10, 1971. The change was the result of a required assessment of existing NAAQS to include a review of new health effects data that have become available since 1970. As a result of this review and national public comments, the standard was changed to a level that is considered to be sufficient to protect the public health and welfare. Since then additional research has concluded that there is in fact damage being caused by ozone levels less than the existing Federal standard. Based on recent studies there appears to be significant vegetation damage at levels considerably below the Federal standard and some "adverse" health effects at the current Federal standard. As of the date this report was compiled no proposals have been made for changing the Federal standard. The current State Standard is .081 ppm. It was established at the same time the original Federal Standard was established and has not been changed. In the past the state standard was interpreted to be .080 ppm but a conversion of the actual 160 ug/m3 standard to ppm yields .081. Therefore, only hourly averages in excess of .081 ppm are considered exceedances of the state standard.

### 3.4 Monitoring

Ozone was monitored at ten sites in Maine during 1987 using continuous monitoring equipment of two kinds, either chemiluminescence or ultra-violet absorption analyzers. Maine's ozone monitoring season is limited to April through October due to the weather conditions which are not conducive to ozone formation at other times of the year.

Table 3-1 is the 1987 Data Summary for Ozone. Table 3-2 presents the Ozone Historical Comparisons and Table 3-3 presents the Ozone Trends.

TABLE 3 - 1  
1987 OZONE DATA SUMMARY  
(Parts Per Million)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>HIGHEST CONCENTRATION</u>	<u>SECOND HIGHEST CONCENTRATION</u>	<u>NUMBER OF VIOLATIONS</u>	
					<u>STATE*</u>	<u>FEDERAL**</u>
<b>ANDROSCOGGIN INTERSTATE AIR QUALITY CONTROL REGION (107)</b>						
Gardiner	Gardiner High School	4879	.119	.112	25	0
Port Clyde	Port Clyde Ozone	3308	.149	.146	82	2
Isle Au Haut	Isle Au Haut Fire Station	3113	.154	.151	87	3
<b>DOWNEAST AIR QUALITY CONTROL REGION (109)</b>						
Acadia National Park	McFarland Hill Ranger Station	8000	.130	.126	44	0
Bedham	Bald Mountain	4771	.125	.117	35	0
<b>METROPOLITAN PORTLAND AIR QUALITY CONTROL REGION (110)</b>						
Cape Elizabeth	Shelter Site	5165	.156	.152	76	3
Cape Neddick	Cape Neddick Light Station	3253	.134	.125	54	1
Kennebunkport	Parson's Way	4168	.152	.145	67	3
<b>NORTHWEST MAINE AIR QUALITY CONTROL REGION (111)</b>						
T4R2 BKP WKR	Sugarloaf Mountain	3398	.093	.092	40	0
T4R2 BKP WKR	Sugarloaf Mountain - Base	2307	.079	.078	0	0

\* Total number of hours minus one greater than .081 ppm.

\*\* Number of days in violation. Not a statistical estimate.

TABLE 3 - 2  
OZONE HISTORICAL COMPARISONS  
(1-Hour Concentrations)

CAPE ELIZABETH  
Shelter Site

<u>YEAR</u>	<u>SECOND HIGH</u>	<u># OF STATE VIOLATIONS</u>
1978	.160 PPM	202
1979	.155 PPM	116
1980	.178 PPM	141
1981	.122 PPM	98
1982	.140 PPM	117
1983	.163 PPM	187
1984	.146 PPM	148
1985	.165 PPM	141
1986	.128 PPM	68
1987	.152 PPM	76

KENNEBUNKPORT  
Parson's Way

<u>YEAR</u>	<u>SECOND HIGH</u>	<u># OF STATE VIOLATIONS</u>
1982	.120 PPM	42
1983	.148 PPM	149
1984	.147 PPM	184
1985	.168 PPM	190
1986	.138 PPM	62
1987	.145 PPM	67

GARDINER  
Gardiner High School

<u>YEAR</u>	<u>SECOND HIGH</u>	<u># OF STATE VIOLATIONS</u>
1980	.117 PPM	54
1981	.122 PPM	31
1982	.120 PPM	56
1983	.140 PPM	99
1984	.112 PPM	89
1985	.133 PPM	84
1986	.110 PPM	17
1987	.112 PPM	25

ACADIA  
McFarland Hill Ranger Station

<u>YEAR</u>	<u>SECOND HIGH</u>	<u># OF STATE VIOLATIONS</u>
1982*	.055 PPM	0
1983	.135 PPM	98
1984	.130 PPM	86
1985	.117 PPM	57
1986	.108 PPM	37
1987	.126 PPM	44

\* Not a complete year.

TABLE 3 - 3  
OZONE TRENDS  
(1-Hour Concentrations)

CAPE ELIZABETH  
Shelter Site

<u>Year</u>	<u>PERCENTILES</u>		
	<u>10%</u>	<u>50%</u>	<u>90%</u>
1978	.015	.035	.065
1979	.018	.036	.070
1980	.019	.035	.065
1981	.015	.032	.056
1982	.018	.036	.058
1983	.018	.034	.061
1984	.019	.040	.064
1985	.022	.038	.062
1986	.016	.033	.055
1987	.018	.035	.055

KENNEBUNKPORT  
Parson's Way

<u>Year</u>	<u>PERCENTILES</u>		
	<u>10%</u>	<u>50%</u>	<u>90%</u>
1983	.008	.027	.058
1984	.012	.032	.064
1985*	.015	.037	.072
1986	.013	.033	.053
1987	.013	.032	.054

\* Percentiles calculated using  
70% of the data.

GARDINER  
Gardiner High School

<u>YEAR</u>	<u>PERCENTILES</u>		
	<u>10%</u>	<u>50%</u>	<u>90%</u>
1980	.008	.031	.056
1981	.009	.029	.050
1982	.009	.030	.053
1983	.009	.031	.056
1984	.007	.031	.055
1985	.012	.034	.057
1986	.009	.029	.047
1987	.008	.028	.048

ACADIA  
McFarland Hill Ranger Station

<u>YEAR</u>	<u>PERCENTILES</u>		
	<u>10%</u>	<u>50%</u>	<u>90%</u>
1982*	.005	.020	.030
1983	.019	.032	.053
1984	.020	.032	.050
1985	.022	.032	.048
1986	.019	.032	.047
1987	.021	.033	.049

\* Not a complete year.

## 4. NITROGEN DIOXIDE (NO<sub>2</sub>)

### 4.1 Description and Sources

In its pure state, nitrogen dioxide is a reddish-orange-brown gas with a characteristic pungent odor. It is corrosive and a strong oxidizing agent. Nitrogen dioxide comprises about 10% of the oxides of nitrogen (NO<sub>x</sub>) that are formed when nitrogen in the air combines with oxygen during high temperature combustion. Most of the rest of the NO<sub>x</sub> emitted by combustion sources is nitric oxide (NO). However, during the day most of the NO is photochemically transformed into NO<sub>2</sub>. Thus, essentially all the NO<sub>x</sub> emitted can be assumed to eventually become NO<sub>2</sub>.

### 4.2 Health and Welfare Effects

Exposure to NO<sub>2</sub> affects the delicate structure of lung tissue. High levels cause lung irritation and potential lung damage. Lower levels have been associated with increased respiratory disease. Oxides of nitrogen can cause serious injury to vegetation, including bleaching or death of plant tissue, loss of leaves, and reduced growth rate. NO<sub>x</sub> also deteriorates fabrics and fades fabric dyes. Nitrate salts formed from nitrogen oxides have been associated with the corrosion of metals. Nitrogen oxides can also reduce visibility.

### 4.3 Standards

The current standard for NO<sub>2</sub> is an annual arithmetic mean (average) value not to exceed .05 ppm. NO<sub>2</sub> is the only gaseous pollutant for which only a long-term (annual average) standard has been established.

### 4.4 Monitoring

Nitrogen Dioxide was monitored at two sites in Maine during 1987. Table 2-1 presents the data collected during 1987.

TABLE 4 - 1  
1987 NITROGEN DIOXIDE DATA SUMMARY  
(Parts Per Million)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>ANNUAL ARITHMETIC MEAN</u>
<b>METROPOLITAN PORTLAND AIR QUALITY CONTROL REGION (110)</b>			
Cape Elizabeth	Shelter Site	3084	.004*
Portland	Shelter Site	571	.049*

\* Insufficient data collected for valid annual arithmetic mean.



## 5. SULFUR DIOXIDE (SO<sub>2</sub>)

### 5.1 Description and Sources

Sulfur dioxide is a colorless irritating gas having the same pungent odor as a struck match. Most people can detect its taste at a level of about 0.3 to 1 part per million. SO<sub>2</sub> is highly soluble in water, forming sulfurous acid. On a worldwide basis, SO<sub>2</sub> is considered to be one of the major pollution problems. It is emitted mainly from stationary sources that utilize fossil fuels (coal, oil) such as power plants, ore smelters, and refineries.

### 5.2 Health and Welfare Effects

The health effects of sulfur dioxide appear to be always associated with high levels of particulates or other pollutants. The world's major recorded air pollution disasters have been associated with high levels of sulfur dioxide and particulates. The excess deaths attributed to these pollutants were due to respiratory failures and occurred predominantly, but not exclusively, in the elderly and infirm. Atmospheres containing high levels of sulfur dioxide are associated with elevated concentrations of other sulfur compounds such as sulfates and sulfuric acid mists, which are corrosive and potentially carcinogenic.

The corrosiveness of SO<sub>2</sub> and its derivatives also causes crop and material damage. Its transport and transformation into sulfurous and sulfuric acids contribute to acid precipitation, causing soils and lakes to become seriously acidified.

### 5.3 Standards

There are two existing Primary National Ambient Air Quality Standards for sulfur dioxide. The first is a long-term one year arithmetic average of 0.03 parts per million (ppm). The second is a short-term 24-hour average standard where concentrations are not to exceed 0.14 ppm more than once per year. The current Secondary NAAQS for SO<sub>2</sub> is a 3-hour average concentration of 0.5 ppm not to be exceeded more than once per year.

In addition there are three state standards for sulfur dioxide. The first is a long-term one-year arithmetic average of .022 parts per million. The second was a short-term 24-hour average standard of .088 ppm not to be exceeded. The third was a short-term 3-hour average concentration of .439 ppm not to be exceeded. During 1987 both of the short-term standards were amended to allow for one exceedance per year.

### 5.4 Monitoring

Sulfur dioxide was monitored at eighteen sites in Maine during 1987 using continuous monitoring equipment utilizing either the pulsed

fluorescent or coulometric methods.

Table 5-1 is the 1987 Data Summary for SO<sub>2</sub>. Tables 5-2 and 5-3 present the SO<sub>2</sub> Historical Comparison Data.

TABLE 5 - 1  
1987 SULFUR DIOXIDE DATA SUMMARY  
(Parts Per Million)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>HIGHEST 3-HOUR AVERAGE</u>	<u>SECOND HIGHEST 3-HOUR AVERAGE</u>	<u>HIGHEST 24-HOUR AVERAGE</u>	<u>SECOND HIGHEST 24-HOUR AVERAGE</u>	<u>ANNUAL ARITH. MEAN</u>
<b>ANDROSCOGGIN INTERSTATE AIR QUALITY CONTROL REGION (107)</b>							
Lewiston	Country Kitchen Parking Lot	6660	.088	.083	.038	.035	.008
Mexico	Hunt's Property	8275	.155	.124	.043	.039	.009
Rumford	Taylor Mountain I	8313	.213	.187	.098	.092	.014
Rumford	Taylor Mountain II	8299	.200	.170	.065	.062	.009
Rumford	Village Green Site	8317	.139	.117	.042	.041	.007
Searsport	Searsport DOT	7732	.173	.171	.056	.040	.004*
<b>AROOSTOOK AIR QUALITY CONTROL REGION (108)</b>							
Madawaska	Madawaska High School	7978	.149	.148	.076	.068	.005
Madawaska	Albert Street	8147	.212	.165	.072	.067	.012
Madawaska	U. S. Post Office	7947	.177	.173	.084	.080	.013
<b>DOWNEAST AIR QUALITY CONTROL REGION (109)</b>							
Bangor	Kenduskeag Pump Station	2967	.095	.092	.050	.039	.009*
East Millinocket	Katahdin School	1072	.164	.122	.054	.030	.006*
Lincoln	Thomas Motel Trailer Park	8082	.079	.079	.039	.029	.004
Lincoln	Fish Hill Base	7784	.075	.040	.018	.015	.002
Lincoln	Fish Hill Peak	8158	.155	.117	.053	.036	.002
Millinocket	Wastewater Treatment Plant	7677	.101	.097	.054	.052	.008
Millinocket	York Street	8507	.247	.151	.048	.045	.007
Woodland	Secondary Treatment Pipeline	7174	.100	.094	.024	.022	.003
<b>METROPOLITAN PORTLAND AIR QUALITY CONTROL REGION (110)</b>							
Portland	Shelter Site	8085	.083	.076	.049	.047	.010

\* Insufficient data collected for valid annual arithmetic mean.

TABLE 5 - 2  
SULFUR DIOXIDE HISTORICAL COMPARISONS  
(Maximum 24-Hour Concentrations of Sulfur Dioxide)

SITE	ADDRESS	MAXIMUM 24-HOUR CONCENTRATION (PPM)					
		1982	1983	1984	1985	1986	1987
ANDROSCOGGIN INTERSTATE AIR QUALITY CONTROL REGION (107)							
Lewiston	Country Kitchen Parking Lot	.056	.044	.060	.043	.047	.038
Mexico	Hunt's Property	--	.061	.071	.070	.068	.043
Rumford	Taylor Mountain I	.075	.077	.096	.066	.086	.098
Rumford	Taylor Mountain II	--	.072	.071	.050	.067	.065
Rumford	Village Green Site	--	.054	.049	.031	.059	.042
Searsport	Searsport DOT	--	--	--	.019	.050	.056
AROOSTOOK AIR QUALITY CONTROL REGION (108)							
Madawaska	Madawaska High School	.139	.049	.066	.037	.046	.076
Madawaska	Albert Street	.152	.130	.078	.058	.080	.072
Madawaska	U. S. Post Office	--	--	--	.061	.068	.084
DOWNEAST AIR QUALITY CONTROL REGION (109)							
East Millinocket	Katahdin School	.072	.054	.025	.026	.027	.054
Lincoln	Thomas Motel Trailer Park	.062	.052	.076	.051	.037	.039
Lincoln	Fish Hill Base	--	.023	.016	.023	.012	.018
Lincoln	Fish Hill Peak	--	.025	.025	.044	.030	.053
Millinocket	Wastewater Treatment Plant	.078	.077	.062	.076	.071	.054
Millinocket	York Street	.063	.065	.044	.046	.061	.048
Woodland	Secondary Treatment Pipeline	.022	.058	.059	.027	.037	.024
METROPOLITAN PORTLAND AIR QUALITY CONTROL REGION (110)							
Portland	Shelter Site	--	.056	.062	.050	.062	.047

TABLE 5 - 3  
SULFUR DIOXIDE HISTORICAL COMPARISONS  
(Sites with violations in past six years)

<u>SITES</u>	<u>ADDRESS</u>	<u>TOTAL NUMBER OF VIOLATIONS*</u>					
		<u>1982</u>	<u>1983</u>	<u>1984</u>	<u>1985</u>	<u>1986</u>	<u>1987</u>
<b>ANDROSCOGGIN INTERSTATE AIR QUALITY CONTROL REGION (107)</b>							
Rumford	Taylor Mountain I	0	0	1	0	0	1
<b>AROOSTOOK AIR QUALITY CONTROL REGION (108)</b>							
Madawaska	Madawaska High School	1	0	0	0	0	0
Madawaska	Albert Street	7	2	0	0	0	0

\* Includes 3-Hour and 24-Hour Violations.

## 6. PARTICULATES (TSP)

### 6.1 Description and Sources

Particulates is the term given to the tiny particles of solid or semi-solid material found in the atmosphere. It is this "dirt" in the air that is visible as a "Brown Cloud", haze or smog. The sources of particulates are many: wind-blown dust and sand from roadways, fields, and construction; coal dust, fly ash, and carbon black from various combustion sources; and automobile exhaust, to name a few. Particulates that range in size from less than 0.1 micrometer up to approximately 45 micrometers are called "total suspended particulates". Particles larger than that range tend to settle out of the air and not remain suspended, except in high winds.

### 6.2 Health and Welfare Effects

The human nose filters out 99 percent of the large and medium-sized particles. The rest enter the windpipe and lungs, where some, known as inhalable particulates, cling to protective mucous and are removed. Some of the smallest, called respirable particulates, are deposited in the lungs' tiny air sacs (alveoli).

In the lungs particulates slow down the exchange of oxygen with carbon dioxide in the blood, causing shortness of breath. The heart may be strained because it must work harder to compensate for oxygen loss. Usually the people most sensitive to these conditions have respiratory diseases like emphysema, bronchitis, asthma, or heart problems. The elderly and children are also sensitive.

Particles themselves may be poisonous if inhaled or absorbed, damaging remote organs like the kidneys or liver. Swallowed mucous that is laden with poisonous particulate matter may damage the stomach.

In addition, particulates may be carriers of poisonous liquid or gaseous substances. Sulfur dioxide, a major air pollutant in its own right, is frequently absorbed by particulates and can react with them to form sulfates. Sulfates react with moisture in the air or in the respiratory tract to form a corrosive liquid (sulfuric acid) that irritates delicate membranes and slows down the cleansing action of mucous. This effect can reduce the body's ability to remove harmful bacteria, increasing the possibility of infection.

Adverse health effects from particulate matter aren't always seen immediately. Particulates can accumulate in the lungs after repeated, long-term exposure, causing respiratory distress and other health problems that may be manifested later.

Particles in the air block out and scatter sunlight, reducing visibility. Particulates soil and corrode metals, masonry, and textiles. Irritating odors are often associated with particulates,

also.

### 6.3 Standards

#### Primary:

At the beginning of 1987 the primary particulate standards were for total suspended particulates (TSP), independent of particle size or chemical composition. The long-term standard was an annual geometric mean not to exceed 75 micrograms of particulates per cubic meter of air ( $\mu\text{g}/\text{m}^3$ ). The short-term standard was a 24-hour average of 260  $\mu\text{g}/\text{m}^3$  not to be exceeded more than once per year.

In July EPA published revised particulate standards to account for the deeper inhalability of smaller particles and eliminated the total suspended particulate standards. The new standards, rather than applying to TSP, apply to inhalable or fine particulates. A particle size of 10 micrometers was selected as the upper size limit with a 24-hour concentration of 150  $\mu\text{g}/\text{m}^3$  and an annual standard of 50  $\mu\text{g}/\text{m}^3$  expressed as an expected annual arithmetic mean (AAM). The short-term standard is attained when the expected number of exceedances is no more than one per year. The expected AAM is determined by averaging the annual arithmetic averages from three successive years of data.

#### Secondary:

The secondary TSP standard was a 24-hour average of 150  $\mu\text{g}/\text{m}^3$  not to be exceeded more than once per year, designed to protect from soiling, corrosion, etc.

When EPA adopted the fine particulate standards they eliminated the secondary TSP standards and made the secondary fine particulate standards equal to the primary fine particulate standards.

#### State Standards:

As of the end of 1987 the State Standards for total suspended particulates included an annual geometric mean of 60 micrograms per cubic meter and a 24-hour standard of 150 micrograms per cubic meter not to be exceeded. The state is expected to adopt the federal fine particulate standards during 1988.

### 6.4 Monitoring

Particulates were monitored at 54 sites in Maine during 1987 using High-Volume Particulate Air Samplers (Hi-Vols).

Hi-Vols operate on the same principle as a vacuum cleaner in that the air is drawn through a filter to "catch the dust". The difference is that a Hi-Vol draws a calibrated volume of air through a pre-weighed filter pad (rather than a bag) for a twenty-four hour period. The change in weight of the filter pad is recorded as total suspended particulate or TSP in micrograms of particulates per cubic meter of air.

Table 6-1 is a summary of the TSP data collected in Maine during 1987. Table 6-2 is a historical comparison of the TSP Annual Geometric Means at sites which have been in existence over the last two years. Table 6-3 summarizes the number of TSP violations which have occurred over the last six years and the sites at which they occurred.

Fine particulate sampling increased again during 1987. The increased sampling was initially conducted to obtain data to evaluate the proposed fine particulate standards and determine those areas which were likely to have problems meeting the proposed range of standards. After July 1, 1987 the sampling was also done to document compliance with the new federal standard. The sampling has been primarily conducted with size-selective hi-vols. One dichotomous sampler was operated at Acadia National Park to support monitoring objectives of the National Park Service. The dichotomous sampler collects particles 10 microns and smaller in two different size classes. The two classes are summed to give total fine particulate. The size-selective hi-vols collect particles 10 microns and smaller. Only the size-selective hi-vols have been approved as federal reference methods for the sampling of fine particulates.

The data collected and the sites which were in operation during 1987 have been summarized in Table 6-4.



TABLE 6 - 1  
1987 TOTAL SUSPENDED PARTICULATES DATA SUMMARY  
(Micrograms Per Cubic Meter)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>HIGHEST 24-HOUR</u>	<u>SECOND HIGHEST</u>	<u>THIRD HIGHEST</u>	<u>ANNUAL GEOMETRIC MEAN</u>
<b>ANDROSCOGGIN INTERSTATE AIR QUALITY CONTROL REGION (107)</b>						
Auburn	Lepage Bakery	60	127	87	87	41.1
Augusta	Hartford Fire House	165	189	185	129	39.6
Jay	Weather Level I	354	175	147	147	34.1
Jay	Crash Road	357	103	87	66	19.4
Jay	Jay Hill	357	118	116	115	25.1
Jay	Burnham	362	131	127	117	33.6
Mexico	Labonville's	223	136	123	118	40.8
Rumford	Taylor Mountain I	226	121	89	82	30.0
Rumford	Taylor Mountain II	225	116	72	66	22.9
Rumford	Village Green Site	226	129	96	91	27.2
Skowhegan	Hinckley	119	93	76	60	18.0
Skowhegan	Eaton Ridge	111	56	47	46	15.5
Thomaston	Mitchell Property	200	156	155	128	21.9
Thomaston	Sander's Property	197	133	96	73	22.6
Thomaston	Pease Property	199	111	101	93	28.4
Thomaston	Marsh Road	197	141	115	97	23.4
Waterville	Stern's Department Store	60	261	117	116	55.1
Winslow	Gulley Hill Road	232	184	155	147	43.6
<b>AROOSTOOK AIR QUALITY CONTROL REGION (108)</b>						
Fort Kent	UMFK-Cyr Hall	9	53	45	34	17.5*
Fort Fairfield	Peterson's	69	117	82	80	33.5*
Madawaska	St. Jarres	116	342	231	197	44.9
Presque Isle	Northeastland Hotel	210	476	334	222	43.8
<b>DOWNEAST AIR QUALITY CONTROL REGION (109)</b>						
Acadia National Park	McFarland Hill Ranger Station	116	44	39	38	12.5
Bangor	Regional Office	58	153	128	92	43.9
Bangor	Kenduskeag Pump Station	114	190	153	135	53.0

TABLE 6 - 1 (continued)  
1986 TOTAL SUSPENDED PARTICULATES DATA SUMMARY  
(Micrograms Per Cubic Meter)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>HIGHEST 24-HOUR</u>	<u>SECOND HIGHEST</u>	<u>THIRD HIGHEST</u>	<u>ANNUAL GEOMETRIC MEAN</u>
Brewer	Brewer Junior High School	57	117	82	75	37.0
Bucksport	Bucks Mill Road	175	64	53	42	12.9*
Bucksport	Public Landing	176	259	181	181	31.5*
East Millinocket	Katahdin School	115	137	123	95	19.2
Lincoln	Vocational Education Building	354	121	120	115	28.8
Lincoln	Lincoln Post Office Building	354	186	153	120	30.3
Lincoln	Thomas Motel Trailer Park	361	143	108	106	33.9
Millinocket	York Street	189	179	176	160	34.4
Old Town	Marsh Island Apartments	61	198	99	99	36.0
Old Town	Penobscot Shoe Company	18	137	131	87	47.4*
Newburgh	Newburgh School	341	69	60	54	15.1
Milford	Shumway Field	322	91	87	83	25.2
Woodland	Secondary Treatment Pipeline	352	245	186	175	24.1
Woodland	Woodland High School	351	181	176	168	29.0
Woodland	Background	344	58	57	52	13.6

**METROPOLITAN PORTLAND AIR QUALITY CONTROL REGION (110)**

Berwick	Berwick Fire Station	117	189	183	167	59.9
Biddeford	Biddeford Treatment Plant	115	110	99	83	36.0
Bridgton	Upper Ridge Road	113	65	58	55	13.0
Brunswick	Cooks Corner	87	143	128	103	43.7*
Portland	Shelter Site	119	160	130	117	48.1
Portland	YWCA	112	102	95	92	43.8
South Portland	SMVTI	116	67	64	59	28.7
Westbrook	N. E. T. & T. Company	112	128	82	81	38.2
Westbrook	Research Building	180	232	227	224	71.2
Westbrook	Warehouse #5	189	172	159	156	60.1

\* Insufficient data collected for valid annual geometric mean.

TABLE 6 - 2  
TOTAL SUSPENDED PARTICULATES HISTORICAL COMPARISON  
ANNUAL GEOMETRIC MEANS (UG/M3)

SITE	ADDRESS	ANNUAL GEOMETRIC MEANS (ug/m3)					
		1982	1983	1984	1985	1986	1987
ANDROSCOGGIN INTERSTATE AIR QUALITY CONTROL REGION (107)							
Auburn	Lepage Bakery	47.4	39.3	43.5	44.8	46.0	41.1
Augusta	Hartford Fire House	46.9	48.1*	45.9	44.3	41.0	39.6
Jay	Weather Level I	40.0	33.0*	36.4	36.6	33.5	34.1
Jay	Crash Road	22.1	18.0*	22.1	18.7	18.9	19.4
Jay	Jay Hill	28.5	25.2*	32.6	24.5	24.6	25.1
Mexico	Labonville's	53.5	50.6	51.6	50.7	46.6	40.8
Rumford	Taylor Mountain I	37.9	34.8	37.5	35.8	33.0	30.0
Rumford	Taylor Mountain II	--	26.0	28.2	26.7	24.3	22.9
Rumford	Village Green Site	--	--	34.0	31.2	29.7	27.2
Skowhegan	Hinckley	18.5	17.3	21.3	18.5	16.6	18.0
Skowhegan	Eaton Ridge	17.4	15.4	20.2	18.4	17.1	15.5
Thomaston	Mitchell Property	25.5	22.0	24.2	22.9	22.0	21.9
Thomaston	Sanders Property	23.7	21.9	25.4	22.9	22.0	22.6
Thomaston	Pease Property	34.0	28.0	31.3	28.4	27.9	28.4
Thomaston	Marsh Road	28.3	22.7	25.9	24.0	23.5	23.4
Waterville	Sterns Department Store	--	--	35.5*	40.8	42.1	55.1
AROOSTOOK AIR QUALITY CONTROL REGION (108)							
Madawaska	St. Jarres	--	--	50.7	46.9	44.7	44.9
Presque Isle	Northeastland Hotel	62.0	66.8	62.5*	59.2	51.6	43.8
DOWNEAST AIR QUALITY CONTROL REGION (109)							
Acadia National Park	McFarland Hill Ranger Station	--	11.6*	12.9	11.6	11.8	12.5
Bangor	Regional Office	42.7	41.7	46.5	44.8	42.3	43.9
Bangor	Kenduskeag Pump Station	52.1	49.8	56.5	59.9	59.4	53.0
Brewer	Brewer Junior High School	36.4	37.0	41.5	38.1	36.5	37.0

TABLE 6 - 2 (continued)  
TOTAL SUSPENDED PARTICULATES HISTORICAL COMPARISON  
ANNUAL GEOMETRIC MEANS (UG/M3)

SITE	ADDRESS	ANNUAL GEOMETRIC MEANS (ug/m3)					
		1982	1983	1984	1985	1986	1987
East Millinocket	Katahdin School	30.8	27.4	25.3	26.9	23.9	19.2
Lincoln	Vocational Education Building	41.5	36.2	35.3	37.1	30.3	28.8
Lincoln	Lincoln Post Office Building	46.6	39.8	40.4	39.2	34.2	30.3
Lincoln	Thomas Motel Trailer Park	44.4	40.9	41.8	41.4	34.9	33.9
Millinocket	York Street	43.3	43.8	49.1	46.1	37.3	34.4
Old Town	Marsh Island Apartments	38.6	35.8	37.3	33.8	32.6	36.0
Old Town	Penobscot Shoe Company	32.1	28.0	31.8	28.1	30.9	47.4
Newburgh	Newburgh School	15.9	15.8	16.1	15.1	16.9	15.1
Milford	Shumway Field	31.6	25.7*	29.1	26.6	25.4	25.2
Woodland	Secondary Treatment Pipeline	31.6*	32.3	--	--	26.5	24.1
Woodland	Woodland High School	36.6*	35.0	--	--	27.3	29.0
Woodland	Background	--	--	--	--	12.0*	13.6

**METROPOLITAN PORTLAND AIR QUALITY CONTROL REGION (110)**

Berwick	Berwick Fire Station	--	--	--	--	51.6*	59.9
Biddeford	Biddeford Treatment Plant	43.0	37.8*	43.3*	35.8	38.8	36.0
Bridgton	Upper Ridge Road	--	--	17.1*	14.6	13.5	13.0
Portland	Shelter Site	48.2	45.6	49.4	51.3	49.7	48.1
South Portland	SMVTI	32.5	33.5*	31.7*	30.7	29.8	28.7
Westbrook	N. E. T. & T. Company	44.0	36.5	40.8	44.7	39.2	38.2
Westbrook	Research Building	55.3	52.2	63.4	70.5	67.4	71.2
Westbrook	Warehouse #5	59.9	51.3	60.6	62.5	57.4	60.1

\* Insufficient data collected for valid annual geometric mean.

TABLE 6 - 3  
TOTAL SUSPENDED PARTICULATES HISTORICAL COMPARISON  
(Sites with violations in past six years)

SITE	ADDRESS	TOTAL NUMBER OF SHORT TERM VIOLATIONS					
		1982	1983	1984	1985	1986	1987
ANDROSCOGGIN INTERSTATE AIR QUALITY CONTROL REGION (107)							
Auburn	Lepage Bakery	4	1	0	0	3	0
Augusta	Hartford Fire House	6	6	18	0	1	2
Jay	Weather Level I	0	1	2	0	0	1
Jay	Crash Road	2	0	0	0	0	0
Jay	Jay Hill	2	0	0	0	0	0
Mexico	Labonville's	0	0	0	1	0	0
Rumford	Village Green	-	1	1	0	0	0
Thomaston	Mitchell Property	1	0	0	0	0	2
Thomaston	Marsh Road	3	0	0	0	0	0
Waterville	Sterns Department Store	-	-	-	-	-	1
Winslow	Gulley Hill Road	-	-	-	-	-	2
AROOSTOOK AIR QUALITY CONTROL REGION (108)							
Madawaska	St. Jarres	-	-	0	1	3	6
Presque Isle	Northeastland Hotel	12	11	12	11	10	6
DOWNEAST AIR QUALITY CONTROL REGION (109)							
Bangor	Regional Office	2	1	0	2	1	1
Bangor	Kenduskeag Pump Station	6	2	1	5	6	2
Bucksport	Public Landing	-	-	-	-	0	4
East Millinocket	Katahdin School	0	1	0	0	0	0
Lincoln	Vocational Education Building	4	2	0	0	0	0
Lincoln	Lincoln Post Office Building	6	7	1	1	1	2
Lincoln	Thomas Motel Trailer Park	10	4	2	3	0	0
Millinocket	York Street	2	3	4	1	1	4
Old Town	Marsh Island Apartments	1	0	2	1	0	1
Old Town	Penobscot Shoe Company	2	0	0	0	0	0
Woodland	Secondary Treatment Pipeline	0	5	1	1	2	3
Woodland	Woodland High School	0	8	11	0	4	5

TABLE 6 - 3 (continued)  
TOTAL SUSPENDED PARTICULATES HISTORICAL COMPARISON  
(Sites with violations in past six years)

<u>SITE</u>	<u>ADDRESS</u>	<u>TOTAL NUMBER OF SHORT TERM VIOLATIONS</u>					
		<u>1982</u>	<u>1983</u>	<u>1984</u>	<u>1985</u>	<u>1986</u>	<u>1987</u>
<b>METROPOLITAN PORTLAND AIR QUALITY CONTROL REGION (110)</b>							
Berwick	Berwick Fire Station	-	-	-	-	1	3
Biddeford	Biddeford Treatment Plant	1	0	0	0	0	0
Portland	Shelter Site	0	0	0	1	0	1
Westbrook	N. E. T. & T. Company	0	0	1	0	0	0
Westbrook	Research Building	4	2	2	8	15	11
Westbrook	Warehouse #5	4	0	1	0	2	4

TABLE 6 - 4  
1987 FINE PARTICULATE DATA SUMMARY  
(Micrograms Per Cubic Meter)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>HIGHEST 24-HOUR</u>	<u>SECOND HIGHEST</u>	<u>THIRD HIGHEST</u>	<u>ANNUAL ARITH. MEAN</u>	<u>ANNUAL GEOM. MEAN</u>
<b>ANDROSCOGGIN INTERSTATE AIR QUALITY CONTROL REGION (107)</b>							
Augusta	Hartford Fire House	150	89	68	67	25.3	22.0
Jay	Jay Hill	115	61	49	49	18.9	15.4
Winslow	Gulley Hill Road	49	73	69	60	28.6	25.3
Mexico	Labonville's	87	107	85	67	30.3	26.0
<b>AROOSTOOK AIR QUALITY CONTROL REGION (108)</b>							
Madawaska	Big Daddy's Restaurant	130	222	143	92	31.8	26.5
Presque Isle	Northeastland Hotel	316	231	172	153	29.2	23.9
<b>DOWNEAST AIR QUALITY CONTROL REGION (109)</b>							
Acadia National Park	McFarland Hill Ranger Sta.(Dichot)	111	41	37	36	12.4	10.5
Hampden	McGraw School	19	39	29	26	15.3	13.1*
Lincoln	Thomas Motel Trailer Park	89	71	63	59	30.8	27.1*
Bangor	Kenduskeag Pump Station	2	24	17	-	20.5	20.2*
Orrington	Center Drive School	20	41	30	21	13.9	12.0*
Woodland	Woodland High School	116	92	73	65	23.4	19.8
<b>METROPOLITAN PORTLAND AIR QUALITY CONTROL REGION (110)</b>							
Bridgton	Upper Ridge Road	40	60	28	27	16.2	14.6
Portland	Shelter Site	50	77	60	58	30.9	28.4
Portland	Elks Lodge	65	55	43	42	21.9	19.7
Westbrook	Warehouse #5	91	94	71	58	28.4	25.9

\* Insufficient data collected for valid annual geometric mean.

## 7. LEAD (Pb)

### 7.1 Description and Sources

Lead in the ambient air exists primarily as particulate matter in the inhalable size range. The predominant source of atmospheric lead is from motor vehicles that burn "leaded" gasoline. The lead in gasoline is in the form of tetraethyl lead, an "anti-knock" compound. Other major sources of atmospheric lead are the extraction and processing of metallic ores.

### 7.2 Health and Welfare Effects

When atmospheric lead is breathed in, it is absorbed into the bloodstream and distributed throughout the body along with lead from contaminated food and drinking water. Lead accumulation in the body can impair the production of hemoglobin. Clinical lead poisoning occurs when the body's accumulation of lead becomes too high. Symptoms of lead poisoning range from loss of appetite, fatigue, cramps and constipation, and pains in the ankles and wrists to loss of power in the arms and legs, anemia, kidney disease, mental retardation, blindness and death. Lead concentrations in the ambient air are not sufficient to produce lead poisoning but they do increase the risk of harm when other sources of lead are present. And, indirectly, lead fallout from automotive exhaust onto soil and street surfaces can be ingested in considerable amounts by infants and young children.

### 7.3 Standards

The current National Ambient Air Quality Standard for lead is a 3-month (calendar quarter) average concentration not to exceed 1.5 micrograms of lead per cubic meter of air.

The current State Standard for lead is a 24-hour average concentration of 1.5 micrograms of lead per cubic meter of air not to be exceeded more than once per year.

### 7.4 Monitoring

Lead was monitored at seven sites in Maine during 1987 by taking samples of the Hi-Vol filters from those sites and analyzing the samples for lead content using an atomic absorption analyzer.

Tables 7-1 and 7-2 are the 1987 Data Summaries for Lead. Table 7-3 presents the Lead Historical Comparison Data.



TABLE 7 - 1  
1987 LEAD DATA SUMMARY  
(Micrograms Per Cubic Meter)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>HIGHEST 24-HOUR</u>	<u>SECOND HIGHEST</u>	<u>THIRD HIGHEST</u>	<u>ANNUAL GEOMETRIC MEAN</u>
<b>ANDROSCOGGIN INTERSTATE AIR QUALITY CONTROL REGION (107)</b>						
Auburn	Lepage Bakery	60	.25	.19	.14	.04
<b>AROOSTOOK AIR QUALITY CONTROL REGION (108)</b>						
Presque Isle	Northeastland Hotel	58	.15	.12	.11	.02
<b>DOWNEAST AIR QUALITY CONTROL REGION (109)</b>						
Bangor	Kenduskeag Pump Station	59	.12	.10	.10	.03
Hampden	McBraw School	19	.03	.02	.01	.01
Orrington	Center Drive School	20	.02	.02	.01	.01
<b>METROPOLITAN PORTLAND AIR QUALITY CONTROL REGION (110)</b>						
Portland	Shelter Site	117	.27	.22	.22	.05
Portland	YWCA	112	.15	.13	.11	.03

TABLE 7 - 2  
1987 LEAD DATA SUMMARY BY QUARTERS  
(Micrograms Per Cubic Meter)

<u>SITE</u>	<u>ADDRESS</u>	<u>1ST</u>	1987 QUARTERLY AVERAGES			<u>4TH</u>
			<u>2ND</u>	<u>3RD</u>		
<b>ANDROSCOGGIN INTERSTATE AIR QUALITY CONTROL REGION (107)</b>						
Auburn	Lepage Bakery	.07	.06	.07		.04
<b>AROOSTOOK AIR QUALITY CONTROL REGION (108)</b>						
Presque Isle	Northeastland Hotel	.08	.04	.02		.03
<b>DOWNEAST AIR QUALITY CONTROL REGION (109)</b>						
Bangor	Kenduskeag Pump Station	.07	.04	.03		.04
Hampden	Mcgraw School	---	---	---		.01
Orrington	Center Drive School	---	---	---		.01
<b>METROPOLITAN PORTLAND AIR QUALITY CONTROL REGION (110)</b>						
Portland	Shelter Site	.12	.07	.05		.06
Portland	YWCA	.07	.04	.04		.04

TABLE 7 - 3  
LEAD HISTORICAL COMPARISONS  
(Micrograms Per Cubic meter)

<u>SITE</u>	<u>ADDRESS</u>	MAXIMUM 24-HOUR CONCENTRATION / AAM					
		<u>1982</u>	<u>1983</u>	<u>1984</u>	<u>1985</u>	<u>1986</u>	<u>1987</u>
ANDROSCOGGIN INTERSTATE AIR QUALITY CONTROL REGION (107)							
Auburn	Lepage Bakery	-----	-----	0.77/0.20	0.40/0.11	0.30/0.07	0.25/0.06
AROOSTOOK AIR QUALITY CONTROL REGION (108)							
Presque Isle	Northeastland Hotel	0.89/0.24	0.93/0.19	0.54/0.13	0.62/0.14	0.20/0.07	0.15/0.04
DOWNEAST AIR QUALITY CONTROL REGION (109)							
Bangor	Kenduskeag Pump Station	0.70/0.24	0.59/0.18	0.53/0.14	0.64/0.15	0.18/0.07	0.12/0.04
METROPOLITAN PORTLAND AIR QUALITY CONTROL REGION (110)							
Portland	Shelter Site	0.91/0.29	0.56/0.20	0.71/0.23	0.53/0.19	0.33/0.11	0.27/0.07

## 8. SULFATES (SO<sub>4</sub>) AND NITRATES (NO<sub>3</sub>)

### 8.1 Description and Sources

Sulfates and Nitrates are compounds of varying harmfulness found everywhere in the atmosphere. They are produced by nature as well as man. Man-made sulfates have their origin in sulfur dioxide while nitrates have theirs in nitrogen oxides. Fine particulate compounds, including sulfates and nitrates are formed from chemical reactions between sulfur dioxide or nitrogen dioxide emitted into the air and other substances present there. These fine particulate compounds have a long atmospheric residence time, can be transported in the air for long distances, and are capable of penetrating deeply into the human respiratory tract.

### 8.2 Health and Welfare Effects

Epidemiological studies of populations exposed to particulate sulfates have shown that atmospheric sulfates, more than sulfur dioxide gas or total suspended particulates, are related to aggravation of asthma, aggravation of heart and lung disease in the elderly, and impairment of lung function in school children. This evidence was obtained from EPA's Community Health and Environmental Surveillance System (CHESS). From these studies, estimates of the sulfate threshold for adverse health effects have been derived, as shown in Table 8-1. However, these epidemiological studies have not been substantiated by laboratory studies.

Both sulfates and nitrates are considered to be contributors to the acid deposition problem.

### 8.3 Standards

There are currently no standards for levels of sulfates in ambient air. EPA is presently working on a standard and is expected to make a proposal in the future.

There are no standards for nitrates nor are there any proposed.

### 8.4 Monitoring

Sulfate levels were measured at six sites in Maine during 1987 by taking samples of the Hi-Vol filters from those sites and analyzing the samples for sulfates using the Automated Technicon II Methylthymol Blue Procedure. There is no standard yet and the monitoring methodology is questionable but the data is being included in this report as an aid to those interested in further information about Maine's air quality. Table 8 - 2 summarizes the sulfate data collected during 1987.

Nitrate levels were measured at six sites in Maine during 1987 by also taking samples of the Hi-Vol filters from those sites and

analyzing the samples using Method 353.1 (Colorimetric, Automated, Hydrazine Reduction). This data, summarized in Table 8 - 3, is also being included in this report as an aid to those interested in further information about Maine's air quality. Nitrate data for 1986 had been reported incorrectly and consequently the table listing that data in the 1986 Annual Report on Air Quality is inaccurate. A corrected table has been printed and is available on request from the Bureau of Air Quality.

TABLE 8-1

SULFATE THRESHOLDS FOR ADVERSE HEALTH EFFECTS

<u>ADVERSE HEALTH EFFECT</u>	<u>THRESHOLD CONCENTRATION FOR SUSPENDED SULFATES</u>
Aggravation of Asthma	6 to 10 Micrograms Per Cubic Meter for 24 Hours.
Aggravation of Heart and Lung Disease in the Elderly	9 Micrograms Per Cubic Meter for 24 Hours
Subtle Decreases in Childhood Lung Function	9 to 13 Micrograms Per Cubic Meter for 1 Year.
Increase in Acute Respiratory Disease in Children	13 Micrograms Per Cubic Meter for 1 Year.

TABLE 8 - 2  
1987 SULFATE DATA SUMMARY  
(Micrograms Per Cubic Meter)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>HIGHEST 24-HOUR</u>	<u>SECOND HIGHEST</u>	<u>THIRD HIGHEST</u>	<u>ANNUAL ARITHMETRIC MEAN</u>
<b>ANDROSCOGGIN INTERSTATE AIR QUALITY CONTROL REGION (107)</b>						
Rumford	Taylor Mountain I	54	31.3	27.0	25.2	11.3
<b>AROOSTOOK AIR QUALITY CONTROL REGION (108)</b>						
Madawaska	St. Jarres	55	21.5	14.5	11.0	6.2
<b>DOWNEAST AIR QUALITY CONTROL REGION (109)</b>						
Acadia National Park	McFarland Hill Ranger Station	114	15.0	13.2	11.0	4.4
Bangor	Regional Office	58	17.5	12.6	10.8	6.8
<b>METROPOLITAN PORTLAND AIR QUALITY CONTROL REGION (110)</b>						
Bridgton	Upper Ridge Road	114	31.9	16.1	13.4	5.4
South Portland	SMVTI	117	19.8	18.0	15.5	6.7

TABLE 8 - 3  
1987 NITRATE DATA SUMMARY  
(Micrograms Per Cubic Meter)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>HIGHEST 24-HOUR</u>	<u>SECOND HIGHEST</u>	<u>THIRD HIGHEST</u>	<u>ANNUAL ARITHMETRIC MEAN</u>
<b>ANDROSCOGGIN INTERSTATE AIR QUALITY CONTROL REGION (107)</b>						
Rumford	Taylor Mountain I	49	7.5	5.0	4.7	1.92
<b>AROOSTOOK AIR QUALITY CONTROL REGION (108)</b>						
Madawaska	St. Jarres	55	3.6	3.1	2.9	1.13
<b>DOWNEAST AIR QUALITY CONTROL REGION (109)</b>						
Acadia National Park	McFarland Hill Ranger Station	113	6.0	5.7	5.4	1.30
Bangor	Regional Office	58	5.2	4.2	4.1	1.60
<b>METROPOLITAN PORTLAND AIR QUALITY CONTROL REGION (110)</b>						
Bridgton	Upper Ridge Road	113	4.9	4.7	4.7	1.39
South Portland	SMVTI	112	6.2	5.7	5.5	2.13

## 9. ATMOSPHERIC DEPOSITION

### 9.1 Description and Sources

As a result of the combustion of tremendous quantities of fossil fuels such as coal and oil, the United States annually discharges approximately 50 million metric tons of sulfur and nitrogen oxides into the atmosphere. Through a series of complex chemical reactions these pollutants can be converted into acids, which may return to earth as components of either rain or snow. This atmospheric deposition, more commonly known as acid rain, may have severe ecological impacts on widespread areas of the environment.

### 9.2 Health and Welfare Effects

While direct health effects from acid rain have not been documented there are numerous indirect effects which could have definite effect on mankind. Atmospheric deposition is known to leach heavy metals such as mercury from rocks causing possible contamination of water supplies. Hundreds of lakes in North America and Scandanavia have become so acidic that they can no longer support fish life. The rain falling on forests and other non-farmlands could, in time, cause extensive changes in the soil chemistry. There is not enough information yet to make it possible to say exactly what the results might be, but there is no reason to think the changes will be beneficial.

### 9.3 Standards

There are no standards in effect or proposed for atmospheric deposition. The only permanent solution to the acid rain problem is to keep the acid levels low. The only practical way of achieving this is by reducing emissions at their sources.

### 9.4 Monitoring

During 1987 there were four sites collecting data on atmospheric deposition. Those four sites included two Bureau maintained sites in Bridgton and Acadia National Park, a University of Maine maintained site in Greenville and a National Weather Service maintained site in Caribou. The samples from these four sites are normally collected every Tuesday morning at 9:00 a.m.. Consequently, the samples are not necessarily a single storm event but are more likely to be a composite of all storm events during the previous week. The samples, if there was a significant storm, are used for field measurements of pH and conductivity and are then packaged up for shipment to the National Atmospheric Deposition Program central laboratory in Illinois. In the central laboratory they are also tested for pH and conductivity as well as additional components. Table 9-1 is a summary of the measurements taken at the central laboratory in Illinois from the DEP operated sites for the year 1987. The sulfate deposition figures were corrected for marine aerosol contribution.



TABLE 9 - 1  
1987 ATMOSPHERIC DEPOSITION DATA SUMMARY

<u>SITE</u>	<u>ADDRESS</u>	<u>MAXIMUM</u>	pH	<u>MEAN</u>	<u>DEPOSITION (Kg/ha)</u>	
			<u>MINIMUM</u>		<u>SO4</u>	<u>NO3</u>
DOWNEAST AIR QUALITY CONTROL REGION (109)						
Acadia National Park	McFarland Hill Ranger Station				Data not available at time of publishing.	
METROPOLITAN PORTLAND AIR QUALITY CONTROL REGION (110)						
Bridgton	Upper Ridge Road				Data not available at time of publishing.	

- \* Precipitation weighted mean.
- \*\* Lab measurements.
- \*\*\* Corrected for marine aerosol and normalized to 52 weeks.

## 10. HYDROCARBONS (HC)

### 10.1 Description and Sources

Hydrocarbons are a class of compounds containing carbon and hydrogen in various combinations. They are found especially in petroleum, natural gas and coal. Some are gaseous, some liquid and some are solid. There are in fact over a thousand hydrocarbon compounds. Many of the polluting hydrocarbons are discharged into the air by incomplete combustion of organic materials. A major source of this kind of hydrocarbon emission is the burning of gasoline in automobiles. Other major contributors are organic solvent evaporation, industrial processes, solid waste disposal and fuel combustion in stationary sources. The control of hydrocarbon emissions are accomplished by combustion process optimization, recovery by mass transfer principles, restriction of evaporative loss and process material and fuel substitution.

### 10.2 Health and Welfare Effects

Hydrocarbon air pollutants enter into and promote the formation of photochemical smog (ozone) and thus contribute to the development of eye irritation and respiratory tract problems. By themselves, hydrocarbons may induce adverse health effects, although there is relatively little quantitative data to relate individual hydrocarbons to the risk of human disease.

### 10.3 Standards

The present State and Federal Standard for non-methane hydrocarbons is a three hour average concentration of 160 ug/m<sup>3</sup>.

### 10.4 Monitoring

Hydrocarbons were not monitored as part of the state's continuous air monitoring program during 1987.