

# MAINE STATE LEGISLATURE



The following document is provided by the  
**LAW AND LEGISLATIVE DIGITAL LIBRARY**  
at the Maine State Law and Legislative Reference Library  
<http://legislature.maine.gov/lawlib>



Reproduced from scanned originals with text recognition applied  
(searchable text may contain some errors and/or omissions)

STATE LAW LIBRARY  
AUGUST, MAINE

# ANNUAL REPORT ON AIR QUALITY 1991



TD  
883.5  
.M2  
M33  
1991

**MAINE**  
DEPARTMENT OF  
ENVIRONMENTAL PROTECTION

DEC 14 1992



**1991 ANNUAL REPORT  
ON AIR QUALITY  
IN THE STATE OF MAINE**

Prepared by  
Jeffrey C. Emery  
Environmental Specialist

Maine Department of Environmental Protection  
Bureau of Air Quality Control  
Division of Field Services

Leighton E. Carver, Division Director  
Dennis L. Keschl, Bureau Director  
Dean C. Marriott, Commissioner



## TABLE OF CONTENTS

	<u>PAGE</u>
TABLE OF CONTENTS	i
LIST OF TABLES	iii
LIST OF FIGURES	v
1. INTRODUCTION	1
1.1 Purpose and Overview	1
1.2 Monitoring Sites	16
1.3 Document Organization	16
1.3.1 Explanation of Data Summary Tables	16
1.3.2 Explanation of Historical Comparison Tables	25
1.3.3 Explanation of Trends Tables	25
2. CARBON MONOXIDE (CO)	26
2.1 Description and Sources	26
2.2 Health and Welfare Effects	26
2.3 Standards	27
2.4 Monitoring	27
3. OZONE (O3)	29
3.1 Description and Sources	29
3.2 Health and Welfare Effects	30
3.3 Standards	30
3.4 Monitoring	30
4. NITROGEN DIOXIDE (NO2)	35
4.1 Description and Sources	35
4.2 Health and Welfare Effects	35
4.3 Standards	35
4.4 Monitoring	35
5. SULFUR DIOXIDE (SO2)	37
5.1 Description and Sources	37
5.2 Health and Welfare Effects	37
5.3 Standards	37
5.4 Monitoring	37
6. PARTICULATES (TSP and PM10)	42
6.1 Description and Sources	42
6.2 Health and Welfare Effects	42
6.3 Standards	43
6.4 Monitoring	43
7. LEAD (Pb)	55
7.1 Description and Sources	55
7.2 Health and Welfare Effects	55
7.3 Standards	55
7.4 Monitoring	55

TABLE OF CONTENTS (continued)

	<u>PAGE</u>
8. SULFATES (SO <sub>4</sub> ) AND NITRATES (NO <sub>3</sub> )	59
8.1 Description and Sources	59
8.2 Health and Welfare Effects	59
8.3 Standards	59
8.4 Monitoring	59
9. ATMOSPHERIC DEPOSITION	62
9.1 Description and Sources	62
9.2 Health and Welfare Effects	62
9.3 Standards	62
9.4 Monitoring	62
10. HYDROCARBONS (HC)	64
10.1 Description and Sources	64
10.2 Health and Welfare Effects	64
10.3 Standards	64
10.4 Monitoring	64

## LIST OF TABLES

<u>TABLE</u>	<u>TITLE</u>	<u>PAGE</u>
1-1	National Ambient Air Quality Standards	2
1-2	State of Maine Ambient Air Quality Standards	3
1-3	Number of Ambient Air Quality Violations by Regions	4
1-4	1991 Ambient Air Quality Monitoring Site Directory	17
2-1	1991 Carbon Monoxide Data Summary	28
3-1	1991 Ozone Data Summary	32
3-2	Ozone Historical Comparisons	33
3-3	Ozone Trends	34
4-1	1991 Nitrogen Dioxide Data Summary	36
5-1	1991 Sulfur Dioxide Data Summary	39
5-2	Sulfur Dioxide Historical Comparisons-Maximum 24-Hour Concentrations of Sulfur Dioxide	40
5-3	Sulfur Dioxide Historical Comparisons-Sites with Exceedances of the Standards in the Past Seven Years	41
6-1	1991 Total Suspended Particulates Data Summary	45
6-2	Total Suspended Particulates Historical Comparison-Annual Geometric Means	47
6-3	Total Suspended Particulates Historical Comparisons-Sites With Samples Greater Than 150 ug/m <sup>3</sup>	49
6-4	1991 Fine Particulate Data Summary	50
6-5	Fine Particulates Historical Comparison-Annual Arithmetic Means	52
6-6	Fine Particulates Historical Comparison-Sites with Samples Greater Than 150 ug/m <sup>3</sup>	54
7-1	1991 Lead Data Summary	56
7-2	1991 Lead Data Summary by Quarters	57
7-3	Lead Historical Comparisons	58



LIST OF TABLES (continued)

<u>TABLE</u>	<u>TITLE</u>	<u>PAGE</u>
8-1	Sulfate Thresholds for Adverse Health Effects	60
8-2	1991 Sulfate Data Summary	61
9-1	1991 Atmospheric Deposition Data Summary	63

LIST OF FIGURES

<u>FIGURE</u>	<u>TITLE</u>	<u>PAGE</u>
1-1	TSP Trends	5
1-2A	PM10 Trends - Southern Maine	7
1-2B	PM10 Trends - Northern Maine	8
1-3A	So. Maine SO2 Trends - 24 Hour	9
1-3B	No. Maine SO2 Trends - 24 Hour	10
1-4A	So. Maine SO2 Trends - AAM	11
1-4B	No. Maine SO2 Trends - AAM	12
1-5	Ozone Trends - Hours of State Violations	13
1-6	Lead Trends - Second High 24 Hour	14
1-7	Lead Trends - AAM	15
1-8	State of Maine, Ambient Air Quality Control Regions	24



## 1. INTRODUCTION

### 1.1 Purpose and Overview

The purpose of this report is to present the air quality monitoring data generated by and for the Maine Department of Environmental Protection, Bureau of Air Quality Control, and to provide a historical perspective from which the significance of that data can be interpreted. Air Quality monitoring measures the concentrations of various pollutants in the ambient air. The monitoring is in response to State and Federal requirements to determine whether the air we breathe is attaining and maintaining National and State Ambient Air Quality Standards which are designed to protect the health and welfare of the public. Federal Primary Standards are intended to protect public health. Federal Secondary Standards are intended to protect public welfare. The State Standards are at least as strict as Federal Standards and in some cases are more strict. The reasoning behind establishing more stringent standards is that generally air quality in Maine is significantly cleaner than in other areas and should remain cleaner. The current Federal and State Standards are presented in Tables 1-1 and 1-2. Table 1-3 is a summary indicating all the violations of ambient air quality standards in the State by regions. Later on in this report those violations will be listed by the sites at which they occurred.

A significant portion of the data collected in the State is collected by industry. The Department has required industry to establish monitoring programs primarily when there are air quality problems associated with the industry, or when an industry is planning to build or expand causing a potential increase in air emissions. The State is still collecting monitoring data for long term trends, special studies and for compliance determinations. Ambient air monitoring by both industry and the State will continue in various regions where necessary until such time as standards are being met and background information has been collected.

Included in this section are some figures which depict some of the results of air quality monitoring and control in the State. Figures 1-1 through 1-7 display trends or the lack of a trend which have been occurring at several long term key sites around the State.

Figure 1-1 depicts the annual geometric means for total suspended particulates at several long term sites. The two highest sites, the Research Building site in Westbrook and the Kenduskeag Pump Station site in Bangor, have shown only minor changes in their annual concentrations of Total Suspended Particulates. With the exception of the site in Lincoln all of the TSP sites included in the graph appear to be showing slight downward trends. The site in Lincoln appears to have reversed the downward trend of earlier years and for the last four years has been showing a gradual increase in concentrations. With the elimination of the total suspended particulate standard the emphasis will be placed on keeping the fine particulates under control but the

**TABLE 1-1  
NATIONAL AMBIENT AIR QUALITY STANDARDS  
(1991)**

<u>Pollutant</u>	<u>Averaging Time</u>	<u>Concentration</u>
Particulates (PM10)	Expected Annual Arithmetic Mean:	
	Primary	50 ug/m3
	Secondary	50 ug/m3
	Twenty-Four Hour:***	
	Primary	150 ug/m3
	Secondary	150 ug/m3
Lead (Pb)	Calendar Quarter	1.5 ug/m3
Carbon Monoxide (CO)	One Hour**	35 ppm
	Eight Hour**	9 ppm
Ozone (O3)	One Hour***	0.12 ppm
Nitrogen Dioxide (NO2)	Annual Arithmetic Mean	0.05 ppm
Sulfur Dioxide (SO2)	Annual Arithmetic Mean	0.03 ppm
	Twenty-Four Hour**	0.14 ppm
	Three-Hour** Secondary	0.50 ppm

\* = Federal Guideline Only.

\*\* = Not to be exceeded more than once per year.

\*\*\* = Statistically estimated number of days with exceedances is not to be more than 1 per year.

ppm = Parts of pollutant per million parts of air.

ug/m3 = Micrograms of pollutant per cubic meter of air.

**TABLE 1-2**  
**STATE OF MAINE AMBIENT AIR QUALITY STANDARDS**  
**(1991)**

<u>Pollutant</u>	<u>Averaging Time</u>	<u>Concentration</u>
Particulates (PM10) (Effective 9-30-89)	Annual Arithmetic Mean	40 ug/m3
	Twenty-Four Hour**	150 ug/m3
Lead (Pb)	Twenty-Four Hour*	1.5 ug/m3
Carbon Monoxide (CO)	One Hour*	35 ppm(40 mg/m3)
	Eight Hour*	9 ppm(10 mg/m3)
Ozone (O3)	One Hour*	.081 ppm(160 ug/m3)
Nitrogen Dioxide (NO2)	Annual Arithmetic Mean	.053 ppm(100 ug/m3)
Sulfur Dioxide (SO2)	Annual Arithmetic Mean	.022 ppm( 57 ug/m3)
	Twenty-Four Hour*	.088 ppm(230 ug/m3)
	Three Hour*	.439 ppm(1150 ug/m3)
Hydrocarbon	Three Hour*	160 ug/m3
Particulates (TSP)***	Twenty-Four Hour	150 ug/m3

\* = Not to be exceeded more than once per year.

\*\* = Statistically estimated number of days with exceedances is not to be more than 1 per year.

\*\*\* = Indication of a nuisance condition only.

PPM = Parts of pollutant per million parts of air.

ug/m3 = Micrograms of pollutant per cubic meter of air.

mg/m3 = Milligrams of pollutant per cubic meter of air.

**TABLE 1-3**  
**NUMBER OF AMBIENT AIR QUALITY VIOLATIONS BY REGIONS**  
**(1991)**



<u>POLLUTANT</u>	<u>REGIONS</u>					<u>TOTALS</u>
	<u>107</u>	<u>108</u>	<u>109</u>	<u>110</u>	<u>111</u>	
<b>Fine Particulate(PM10)</b>						
Annual Arithmetic Mean						
State	0	0	0	0	?	0
Federal	0	0	0	0	?	0
Twenty-four Hour						
State	0	0	0	0	?	0
Federal	0	0	0	0	?	0
<b>Lead</b>						
Twenty-four Hour						
State	0	?	0	0	?	0
Federal	0	?	0	0	?	0
<b>Carbon Monoxide</b>						
One Hour	?	?	0	?	?	0
Eight Hour	?	?	0	?	?	0
<b>Ozone</b>						
One Hour						
State	341	?	183	305	?	829
Days						
Federal	10	?	2	11	?	23
<b>Nitrogen Dioxide</b>						
Annual Arithmetic Mean	?	?	?	0	?	0
<b>Sulfur Dioxide</b>						
Annual Arithmetic Mean						
State	0	0	0	0	?	0
Federal	0	0	0	0	?	0
Twenty-four Hour						
State	0	0	0	0	?	0
Federal	0	0	0	0	?	0
Three Hour						
State	0	0	0	0	?	0
Federal	0	0	0	0	?	0

\* Annual Means generated by only a few samples are not included in this summary.

? No monitoring done for this pollutant within this region during 1991.

# FIGURE 1-1

## TSP TRENDS





TSP will be tracked to ensure that any source emissions are properly controlled.

Figures 1-2A and 1-2B indicate trends over the last seven years in the annual arithmetic mean for fine particulate. The majority of the sites collecting PM10 data over the last seven years are showing either a downward trend or are low enough that they are probably recording regional background concentrations and are not indicating a significant trend in either direction. One exception appears to be the site in Madawaska which is recording concentrations in the 30-40 microgram range and is remaining relatively constant. These levels are probably due to a regional background level plus a relatively constant contribution from the winter sanding of the streets in Madawaska. The KPS site in Bangor had shown a significant reduction for the first three years but 1991 reversed the trend. The levels are low enough that unless the trend continues there is little concern at this point



Figures 1-3A, 1-3B, 1-4A and 1-4B indicate the sulfur dioxide trends at seven sites with a long term history. The twenty-four hour concentrations at all the sites indicated showed a decrease in 1991, with the exception of Portland which showed a slight increase. The maximum twenty-four hour concentration, which occurred in Madawaska, was approximately 82% of the standard. The next highest twenty-four hour concentration, which isn't included on the graph, was only 60% of the standard. The annual arithmetic means at all sites are well below the standard with the maximum value indicated for 1991 being one third of the standard. The only site showing an increase during 1991 in the annual concentrations was the Albert Street site in Madawaska. Even with the increase the Madawaska site has still shown a significant downward trend over the last ten years. With the exception of upset or unusual conditions at a large emission source there doesn't appear to be any problems with the sulfur dioxide standards.

Figure 1-5 depicts the number of hourly violations of the State ozone standard. As can be seen from the graphs, the violations vary greatly from year to year and while showing a very significant increase in the number of violations during 1988 there was a significant decrease during 1989 which has continued through 1991 at most of the sites. Weather conditions are responsible for a lot of the variability from year to year and the conditions during 1988 were very conducive to the formation of ozone while those of subsequent years were not. Because of the significant effect weather has on the formation of ozone, Maine, as well as the rest of the northeast, will need to control emissions to such a level that even under ideal weather conditions ozone levels can be kept below the standards.

Figures 1-6 and 1-7 indicate the very significant reduction that has occurred in lead levels throughout the state in both short term concentrations and in the annual arithmetic means. These significant downward trends are primarily due to the decreased use of lead in gasoline. Current lead levels are less than 20% of the state standard and even less of the federal standard and are expected to remain at those levels with only minor fluctuations expected in either direction.



# FIGURE 1-2A

## PM10 TRENDS - SOUTHERN MAINE



# FIGURE 1-2B



## PM10 TRENDS - NORTHERN MAINE



# FIGURE 1-3A


## SO. MAINE SO2 TRENDS - 24 HOUR

Page 9



# FIGURE 1-3B



## NO. MAINE SO2 TRENDS - 24 HOUR



—x— MADAWASKA —■— MILLINOCKET —□— LINCOLN


# FIGURE 1-4A

## SO. MAINE SO2 TRENDS - AAM



# FIGURE 1-4B



## NO. MAINE SO2 TRENDS - AAM



—x— MADAWASKA    —■— MILLINOCKET    —□— LINCOLN

# FIGURE 1-5


## OZONE TRENDS - HOURS OF STATE VIOLATION






# FIGURE 1-6

## LEAD TRENDS - SECOND HIGH 24 HOUR



—x— PORTLAND —■— BANGOR

**FIGURE 1-7**  
**LEAD HISTORICAL TRENDS - AAM**



—x— Portland —■— Bangor

Data summarized in this report is available for review in the Department headquarters in Augusta and copies can be obtained from that office for a nominal fee.

## 1.2 Monitoring Sites

Air quality data are developed using two basic methods; 1) the continuous monitoring of gaseous pollutants and; 2) the periodic sampling of particulate and gaseous pollutants. In addition to pollutant monitoring there is also the continuous monitoring of meteorological parameters.

Continuous gaseous monitoring was done at thirty sites in Maine during 1991. Carbon Monoxide was monitored at one of these stations, ozone at eleven, nitrogen dioxide at one and sulfur dioxide at nineteen.

Particulate sampling was done at forty-six sites in Maine during 1991. Twenty-five of these stations monitored total suspended particulates. Thirty-nine of these sites also collected fine particulate fractions. Also, lead monitoring was done at eight stations. Four sites were analyzed for sulfates. There were also two sites collecting acid rain data as part of the state monitoring network.

In addition to pollutant monitoring, wind speed and direction was recorded at twenty-seven sites around the State during 1991. Some of these sites also recorded other meteorological parameters such as sigma (stability) and temperature, precipitation and solar radiation.

Table 1-4 presents all the monitoring sites in Maine that operated during 1991 and indicates which parameters were monitored at each site. The map in Figure 1-8 shows the Air Quality Control Regions within the State.

## 1.3 Document Organization

This document is divided by pollutant into chapters. Each chapter contains: 1) a description of the nature and sources of that pollutant, 2) its health and welfare effects, 3) a discussion on the standards (current and proposed) for that pollutant, 4) a discussion of the monitoring methods for that pollutant, 5) a table presenting the 1991 monitored data, 6) in the case of some pollutants, historical tables presenting 1991 data along with data for previous years to show trends, effects of control strategy, or change in emission sources.

### 1.3.1 Explanation of Data Summary Tables

The Data Summary Tables were designed to facilitate comparing 1991 air quality monitoring data with the standards for each pollutant. Therefore, the data are presented for each averaging time for which standards exist for a pollutant.

An annual average concentration is presented for each pollutant

TABLE 1-4  
1991 AMBIENT AIR MONITORING SITE DIRECTORY

<u>SITE</u>	<u>ADDRESS</u>	<u>OPERATOR</u>	<u>PARAMETERS MEASURED</u>
ANDROSCOGGIN COUNTY			
Auburn (23 001 0005)	Lewiston-Auburn Airport Lewiston Junction Road	DEP	WS/WD
Lewiston (23 001 0011)	Country Kitchen Parking Lot Canal Street	DEP	SO <sub>2</sub> , TSP, Pb, FP
Livermore Falls (23 001 0013)	James River/Otis Mill Route 4	James River Corporation	WS/WD, Temperature, SO <sub>2</sub> (n)
Livermore Falls(NEW) (23 001 2002)	10-12 Millett Street	Pine State Power	SO <sub>2</sub>
AROOSTOOK COUNTY			
Madawaska (23 003 0006)	Fraser Paper Company Bridge Street	Fraser Paper	WS/WD, Temperature
Madawaska (23 003 0009)	Albert Street	Fraser Paper	SO <sub>2</sub> , Precipitation
Madawaska (23 003 0012)	U. S. Post Office 430 E. Main Street	Fraser Paper	SO <sub>2</sub> , WS/WD
Madawaska (23 003 0013)	Big Daddy's Restaurant 395 E. Main Street	DEP	FP
Madawaska (23 003 1003)	Madawaska High School 7th Avenue	Fraser Paper	SO <sub>2</sub>
Presque Isle (23 003 1005)	Northeastland Hotel 436 Main Street	DEP	FP
Presque Isle (23 003 1008)	Regional Office 528 Central Drive	DEP	WS/WD, FP
T12RB(DISC) (23 003 6001)	Bald Mountain Project	Boliden Resources, Inc.	FP, WS/WD, Temperature
CUMBERLAND COUNTY			
Bridgton (23 005 0002)	Upper Ridge Road	DEP	Acid Precipitation, Sulfate, Nitrate, FP, Precipitation

TABLE 1-4(Continued)  
1991 AMBIENT AIR MONITORING SITE DIRECTORY

<u>SITE</u>	<u>ADDRESS</u>	<u>OPERATOR</u>	<u>PARAMETERS MEASURED</u>
Portland (23 005 0010)	Chevrus High School Ocean Avenue	DEP	WS/WD
Portland (23 005 0014)	Shelter Site(P.E.O.P.L.) Elm Street	DEP	SO2,Pb,FP,Sulfate, NO2,NO,NOX
Portland (23 005 0015)	Tukey's Bridge	DEP	Pb,FP(n)
Westbrook (23 005 1008)	Research Building S. D. Warren	S. D. Warren	TSP,FP
Westbrook (23 005 1009)	S. D. Warren Company Wind S. D. Warren Property	S. D. Warren	WS/WD,Temperature
Cape Elizabeth (23 005 2003)	Shelter Site Two Lights State Park	DEP	Ozone(s),WS/WD(s)
South Portland (23 005 6002)	SMVTI Vocational Drive	DEP	Sulfate
Westbrook (23 005 7002)	N. E. T.& T. Company Ash Street	S. D. Warren	FP
FRANKLIN COUNTY			
Jay (23 007 0003)	Crash Road Gilbert Jewell Property	International Paper	TSP
Jay (23 007 0004)	Jay Hill Bomaster Property	International Paper	TSP,FP
Jay (23 007 0008)	Burnham Site	International Paper	TSP
Jay (23 007 2001)	Weather Level I Lagoon Hill	International Paper	TSP,FP,WS/WD,Temperature, Solar Radiation,Precipitation
HANCOCK COUNTY			
Acadia National Park (23 009 0003)	McFarland Hill Ranger Station Route #233	NPS/DEP	Acid Precipitation, Precipitation
Acadia NP (23 009 0101)	Acadia NP Route #233	NPS	Ozone,SO2(d),WS/WD(n), Temperature(n),Dewpoint(n)

TABLE 1-4(Continued)  
1991 AMBIENT AIR MONITORING SITE DIRECTORY

<u>SITE</u>	<u>ADDRESS</u>	<u>OPERATOR</u>	<u>PARAMETERS MEASURED</u>
Bucksport (23 009 1005)	Waste Disposal Site Route #15	Champion International	WS/WD, Temperature, Precipitation
Bucksport(DISC) (23 009 1006)	Napa Auto Parts 240 Main Street	Applied Energy Services, Inc.	SO2
Dedham (23 009 2003)	Bald Mountain	DEP	Ozone(s), WS/WD(s), CO(d)
KENNEBEC COUNTY			
Augusta (23 011 0008)	Governor's Hangar State Airport	DEP	WS/WD
Augusta(NEW) (23 011 0014)	Rines Hill Parking Lot Water Street	DEP	FP
Waterville (23 011 1004)	Front Street Municipal Park	DEP	TSP, Fp, SO2
Winslow (23 011 2003)	Gulley Hill Road	Scott Paper Company	TSP
Winslow (23 011 2004)	Boston Avenue	Scott Paper Company	TSP, FP
Gardiner(NEW) (23 011 2005)	Pray Street School	DEP	Ozone(s)
Augusta(NEW) (23 011 2007)	DEP Response Building AMHI Grounds	DEP	Ozone(s)
KNOX COUNTY			
Isle Au Haut (23 013 0003)	Isle Au Haut Fire Station	UM/DEP	Ozone(s)
Port Clyde (23 013 0004)	Port Clyde Ozone St. George	DEP	Ozone(s)
Thomaston (23 013 1005)	Dragon Cement Weather Route #1	Dragon Products	WS/WD
Thomaston (23 013 1007)	Marsh Road	Dragon Products	TSP, FP

TABLE 1-4(Continued)  
1991 AMBIENT AIR MONITORING SITE DIRECTORY

<u>SITE</u>	<u>ADDRESS</u>	<u>OPERATOR</u>	<u>PARAMETERS MEASURED</u>
Thomaston (23 013 2001)	Mitchell Property 2 Dexter Avenue	Dragon Products	TSP,FP
OXFORD COUNTY			
Mexico (23 017 0008)	Labonville's Route #2	Boise Cascade	TSP,FP
Mexico (23 017 0011)	Hunt's Property Route #2	Boise Cascade	SO2
Rumford (23 017 2002)	Boise Cascade Weather II Swift River Pump House	Boise Cascade	WS/WD, Temperature, Solar Radiation
Rumford (23 017 2005)	Taylor Mountain I	Boise Cascade	TSP,SO2,Sulfate,FP(n)
Rumford (23 017 2006)	Taylor Mountain II	Boise Cascade	SO2
Rumford (23 017 2007)	Village Green Site	Boise Cascade	TSP,SO2,FP
PENOBSCOT COUNTY			
Bangor (23 019 0002)	Kenduskeag Pump Station Washington Street	DEP	TSP,Pb,FP
Bangor (23 019 0010)	BIA-Building #489 Air National Guard	DEP	WS/WD
Brewer (23 019 1002)	Brewer Junior High School 5 Somerset Street	DEP	TSP,FP
Lincoln (23 019 1007)	Thomas Motel Trailer Park 39 West Broadway	Lincoln Pulp & Paper Company	TSP,SO2,FP
Lincoln(DISC) (23 019 1010)	Lincoln Airport	Lincoln Pulp & Paper Company	WS/WD
Lincoln (23 019 1011)	Lincoln Street	Lincoln Pulp & Paper Company	FP
Lincoln (23 019 1012)	Penobscot River	Lincoln Pulp & Paper Company	FP

TABLE 1-4(Continued)  
1991 AMBIENT AIR MONITORING SITE DIRECTORY

<u>SITE</u>	<u>ADDRESS</u>	<u>OPERATOR</u>	<u>PARAMETERS MEASURED</u>
Lincoln(NEW) (23 019 1013)	Lincoln Mill Katahdin Avenue	Lincoln Pulp & Paper Company	WS/WD, Temperature
Lincoln (23 019 2003)	Lincoln Post Office Building 50 Fleming Street	Lincoln Pulp & Paper Company	TSP, FP
Millinocket(DISC) (23 019 2007)	Katahdin Nursing Home	Great Northern Paper Company	FP
Millinocket (23 019 2009)	York Street	Great Northern Paper Company	TSP, SO2, FP
East Millinocket(DISC) (23 019 2011)	Library/Municipal Building 53 Main Street	Great Northern Paper Company	FP
East Millinocket(DISC) (23 019 2012)	Mill Entrance Main Street	Great Northern Paper Company	SO2
Millinocket (23 019 2013)	Mill Stone Dam	Great Northern Paper Company	WS/WD, Temperature
Old Town (23 019 4003)	Marsh Island Apartments 100 South Main Street	DEP	TSP, FP
Milford (23 019 4006)	Costigan French Settlement Road	DEP	Ozone(s)
Old Town(NEW) (23 019 5004)	Woodlands Garage James River Corporation	James River Corporation	WS/WD, Temperature
Orrington (23 019 8001)	Center Drive School	Penobscot Energy Recovery Co.	FP
Hampden (23 019 8011)	McGraw School	Penobscot Energy Recovery Co.	FP
PISCATAQUIS COUNTY			
Greenville (23 021 0001)	Squaw Brook Greenville	University of Maine	Acid Precipitation, Precipitation
SAGadahoc COUNTY			
Bath (23 023 0002)	Coal Pocket Site	Bath Iron Works	SO2, FP, WS/WD



TABLE 1-4(Continued)  
1991 AMBIENT AIR MONITORING SITE DIRECTORY

<u>SITE</u>	<u>ADDRESS</u>	<u>OPERATOR</u>	<u>PARAMETERS MEASURED</u>
SOMERSET COUNTY			
Madison (23 025 1004)	The Ballfield Main Street	Madison Paper Industries	WS/WD, Temperature
Skowhegan (23 025 2001)	Hinckley Hinckley Farm School	S. D. Warren	TSP, FP
Skowhegan (23 025 2002)	Eaton Ridge	S. D. Warren	TSP, FP
Skowhegan (23 025 2003)	Somerset Mill S. D. Warren	S. D. Warren	WS/WD, Ozone(n)
WASHINGTON COUNTY			
Woodland (23 029 0007)	Secondary Treatment Pipeline	Georgia Pacific Corporation	FP
Woodland (23 029 0008)	Woodland High School	Georgia Pacific Corporation	TSP, FP
Woodland (23 029 0017)	Woodyard Woodland Mill	Georgia Pacific Corporation	WS/WD
Woodland (23 029 0018)	Background	Georgia Pacific Corporation	FP
Jonesport (23 029 0019)	Public Landing	DEP	Ozone(s)
Woodland (23 029 0020)	100 Meter Tower	Georgia Pacific Corporation	WS/WD, Temperature, Dewpoint
YORK COUNTY			
Biddeford (23 031 0002)	Biddeford Treatment Plant Water Street	DEP	TSP, Pb, SO2, FP
Biddeford (23 031 0004)	Biddeford Rotary Park	DEP	WS/WD
Biddeford (23 031 0005)	Eagles Aerie 57 Birch Street	DEP	TSP, Pb

TABLE 1-4(Continued)  
1991 AMBIENT AIR MONITORING SITE DIRECTORY

<u>SITE</u>	<u>ADDRESS</u>	<u>OPERATOR</u>	<u>PARAMETERS MEASURED</u>
Saco (23 031 0006)	Saco Island - CMP	DEP	Pb
Saco (23 031 0007)	Ames Store Roof Spring Street	DEP	Pb
Kennebunkport (23 031 2002)	Parson's Way	DEP	Ozone(s)
Isle of Shoals(NEW) (23 031 4753)	Appledore Island	EPA	Ozone(s)

(AIRS Site #)	New - Site established in 1991	n - Instrument installed during 1991
	DISC - Site discontinued in 1991	d - Instrument removed during 1991
	TSP - Total Suspended Particulate	s - Instrument operated seasonally during 1991
	SO2 - Sulfur Dioxide	i - Instrument operated intermittently during 1991
	NO - Nitric Oxide	
	NOX - Oxides of Nitrogen	
	NO2 - Nitrogen Dioxide	
	CO - Carbon Monoxide	
	Pb - Lead	
	WS/WD - Wind Speed and Direction	
	FP - Fine Particulate	
	NMHC - Nonmethane Hydrocarbons	



Northwest Maine  
Air Quality Control  
Region (111)

Aroostook  
Air Quality Control  
Region (108)

Downeast Air  
Quality Control  
Region (109)


Androscoggin Interstate  
Air Quality Control Region (107)

Metropolitan Portland  
Air Quality Control Region (110)

MINOR CIVIL DIVISIONS

STATE OF  
**MAINE**  
PREPARED BY THE  
STATE OF MAINE  
DEPARTMENT OF TRANSPORTATION  
BUREAU OF PLANNING  
IN COOPERATION WITH THE  
U.S. DEPARTMENT OF TRANSPORTATION  
FEDERAL HIGHWAY ADMINISTRATION

FIGURE 1-8: AIR QUALITY CONTROL REGIONS  
IN THE STATE OF MAINE



that has a long-term, annual standard (NO<sub>2</sub>, SO<sub>2</sub>, PM<sub>10</sub>).

For pollutants that have short-term standards, the highest short-term values are presented. Some pollutants are allowed to exceed the standard once during the year so the second highest value would be used to determine whether there was a violation or not.

All of the data collected during 1991 has been presented in the Data Summary Tables. However, in making comparisons of the data, one should be aware that a site with only a few samples will not be a valid indicator of pollutant concentrations in the area.

### 1.3.2 Explanation of Historical Comparison Tables

The Historical Comparison Tables present air quality data for 1991 and those years prior to 1991 when the same pollutant was monitored at the same site. The purpose of the Tables are to indicate the variations in air quality from year to year. The Tables in some cases represent maximum concentrations for specific time periods and in others the number of days in each year that the standards were violated.

### 1.3.3 Explanation of Trends Tables

The highest hourly concentration in a year is not the best indicator of long-term air quality trends because it is an erratic value. Therefore, special trend tables are presented for carbon monoxide and ozone. The trend tables present the 10th, 50th, and 90th percentile values to represent the bulk of the air quality data for each year. Percentiles indicate the fraction, or percent, of the value that are below a particular level. For example, if the 90th percentile value for some set of CO observations is 5.0 ppm, it means that 90% of the time the concentrations of CO are less than 5.0 ppm. Conversely, it also means that 10% of the time the concentrations are above 5.0 ppm. Thus the existence or lack of long-term trends in overall air quality for CO and O<sub>3</sub> can be more reliably determined using the Trends Tables, than by looking at just the Historical Comparison Tables.

The ozone data has been incorrectly interpreted for the last three years and consequently the trends tables have not reported the ozone data in a consistent format. Starting in 1988 data was stored in the AIRS database instead of the old SAROAD system. Under the SAROAD system reports were generated which indicated the percentiles or frequency distribution of all the hourly data values reported. Reports generated under the AIRS system are now based on percentiles calculated based on each days maximum hourly value. Thus, a 50th percentile of .050 ppm means that 50% of the days monitored had a maximum hourly value during the day of .050 ppm or less. All of the trend information for ozone in this report have been revised to reflect this method of calculation based on the AIRS report format. If a report format can be developed that will provide percentiles based on total hourly values then that statistic will be used in future reports as it is a better indicator of total exposure to high ozone levels.

## 2. CARBON MONOXIDE (CO)

### 2.1 Description and Sources

Carbon monoxide is colorless, odorless and tasteless gas. Therefore you do not even know you are breathing it until you feel its detrimental effects. It constitutes the largest single fraction of the pollutants found in urban atmospheres. It is produced primarily by the incomplete combustion of organic materials used as fuels for transportation and in the heating of buildings; it also results from industrial processes, refuse burning, and agricultural burning. Several natural sources of CO of both biological and non-biological origin have also been identified, but their contributions to urban atmospheric concentrations are thought to be small. Background levels of CO (resulting from natural and technological sources) found in relatively nonpolluted air range from 0.025 to 1.0 ppm. Urban carbon monoxide is produced primarily by motor vehicles.

Because motor vehicle traffic is the major source of CO, daily concentration peaks coincide with morning and evening rush hours. The worst carbon monoxide problems are found where large numbers of slow moving cars congregate. These problems are further aggravated when they occur in a "street canyon" situation. When there are large amounts of slow moving traffic in a street canyon situation, with the wind blowing perpendicular to the street, carbon monoxide can be trapped in the canyon and build up to unhealthy levels.

CO problems are usually worse in winter because: 1) cold weather makes motor vehicles run dirtier and requires more combustion for space heating; and 2) on winter nights a strong inversion layer develops in the atmosphere, that traps pollution near the ground, preventing it from mixing with cleaner air above.

### 2.2 Health and Welfare Effects

Carbon monoxide affects the central nervous system by depriving the body of the oxygen it needs. Tests of automobile drivers show exposure to carbon monoxide can impair driver's judgement and ability to respond rapidly in traffic. It can also impair vision and produce headaches.

Carbon monoxide enters the bloodstream by combining with hemoglobin, the substance that carries oxygen to the cells. Hemoglobin that is bound up with CO is called carboxyhemoglobin. This combination occurs 200 times more readily with CO than with oxygen, so the amount of oxygen being distributed throughout the body by the bloodstream is reduced in CO's presence. Blood laden with CO can weaken heart contractions, lowering the volume of blood distributed to various parts of the body. It can also significantly reduce a healthy person's ability to perform manual tasks, such as working, jogging and walking. A life-threatening situation exists in patients with heart disease, who can't compensate for the oxygen loss. The 4.2 million people in the U.S. suffering from angina pectoris (a heart disease characterized by

brief spasmodic attacks of chest pain due to insufficient oxygen levels in the heart muscles) are especially susceptible. Carbon monoxide is also harmful to persons who have lung disease, anemia or cerebral-vascular disease. Others sensitive to carbon monoxide include the human fetus, and people exposed to long-term concentrations, such as traffic officers.

People who sit in idling cars over sustained periods risk harmful CO exposure, as do cigarette smokers. Since about two percent of cigarette smoke is carbon monoxide, if you or someone else smokes while driving in heavy traffic, you may both experience the harmful effects of CO from the cigarette smoke and the engine exhaust accumulated in streets. Even three or four hours after you're exposed, half the excess CO still remains in your bloodstream. Because it takes time for CO to build up in the bloodstream, the severity of health effects depends both on the concentration being breathed and the length of time the person is exposed.

### 2.3 Standards

The existing standards for carbon monoxide are currently set at 9 parts CO per million parts air (ppm), averaged over a period of 8 hours, and 35 ppm averaged over 1 hour, not to be exceeded more than once per year. As a result of a review and revision of the health criteria, EPA had proposed to retain the existing primary 8-hour standard at 9 ppm and to lower the primary 1-hour standard to 25 ppm. However, this change has not occurred and the standards remain the same. The change in the 1-hour standard had been proposed because of the more rapid accumulation of blood carboxyhemoglobin in moderately exercising sensitive persons compared to resting individuals. The impact of exercise, which is greater for short-duration exposures, was not considered in the original standard.

### 2.4 Monitoring

Carbon monoxide was monitored at one site in Maine during 1991 using continuous monitoring equipment utilizing the non-dispersive infrared technique. Table 2-1 is the 1991 Data Summary for CO. This data was collected primarily to determine existing background levels of carbon monoxide at a rural location.

TABLE 2 - 1  
 1991 CARBON MONOXIDE DATA SUMMARY  
 (Parts Per Million)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>1-HOUR CONCENTRATIONS</u>		<u>8-HOUR CONCENTRATIONS</u>		<u>ANNUAL ARITH. MEAN</u>
			<u>HIGHEST</u>	<u>SECOND HIGHEST</u>	<u>HIGHEST</u>	<u>SECOND HIGHEST</u>	
HANCOCK COUNTY Dedham	Bald Mountain	737	1.0	1.0	1.0	1.0	0.41*

\* Insufficient data for a valid annual arithmetic mean.

### 3. OZONE (O3)

#### 3.1 Description and Sources

Ozone is a highly reactive form of oxygen which, at very high concentrations, is a blue unstable gas that has a characteristic pungent odor most commonly identified around an arcing electric motor, lightning storms, or other electrical discharges. However, at normal ambient concentrations, ozone is colorless and odorless. Ozone is the major component of photochemical "smog", but the haziness and odors of smog are primarily caused by other components.

Natural ground level ozone occurs in low concentrations (less than .05 ppm) due to natural physical and chemical phenomena. Occasionally, unique meteorological conditions can result in natural levels between .05 and .10 ppm.

Ozone is not emitted directly from a source as are other pollutants. It forms as secondary pollutant. Its precursors are hydrocarbons and nitrogen oxides, which chemically react in sunlight to form ozone. The hydrocarbons are emitted in automobile exhaust, from gasoline and oil storage and transfer, and from industrial use of paint solvents, degreasing agents, cleaning fluids, ink solvents, incompletely burned coal or wood and many other sources. Plants also give off hydrocarbons such as terpenes from pine trees. Nitrogen oxides are emitted by all combustion sources.

The highest ozone levels generally occur during summer afternoons when the high temperatures and strong sunlight promote photochemical reactions. Stagnant weather may cause smog to remain in an area for several days. The winds may also transport ozone many miles outside of the urban environment. For example, it is estimated that the majority of the ozone in the State of Maine is transported into the State from sources located outside the State. In addition a much smaller amount of the ozone is naturally occurring background concentrations, part of which is also transported into the State. The remaining ozone is assumed to be due to local sources within the State. Because of long-range transport, local control of emissions by itself may not solve the ozone problem. An effective national program may be necessary to achieve national compliance.

Ground-level ozone, discussed above, should not be confused with the stratospheric ozone layer, located about seven miles high in the atmosphere, which shields the earth from cancer-causing ultraviolet rays. Concentrations of ozone in this layer may reach as high as 10 ppm. Concern over potential reduction of the necessary levels of ozone in the stratosphere by reactions with fluorocarbons from aerosol cans has resulted in the removal of most of these propellants from the market. However, ozone at ground level, where it is breathed, is a pollutant.



### 3.2 Health and Welfare Effects

Ozone at low concentrations causes eye irritations and at higher concentrations difficulty in breathing for people with respiratory problems, the elderly, and children. Many plants, such as white pine, soybeans and alfalfa, are extremely sensitive to ozone, and ozone is known to weaken materials such as rubber and fabrics.

### 3.3 Standards

The existing National Ambient Air Quality Standard (NAAQS) for ozone is 0.12 ppm and will be attained when "the expected number of days per calendar year with maximum hourly average concentrations above 0.12 ppm is equal to or less than one". This standard was effective February 8, 1979 and replaced a more restrictive 0.08 ppm standard that was established April 10, 1971. The change was the result of a required assessment of existing NAAQS to include a review of new health effects data that have become available since 1970. As a result of this review and national public comments, the standard was changed to a level that is considered to be sufficient to protect the public health and welfare. Since then additional research has concluded that there is in fact damage being caused by ozone levels less than the existing Federal standard. Based on recent studies there appears to be significant vegetation damage at levels considerably below the Federal standard and some "adverse" health effects at the current Federal standard. As of the date this report was compiled no proposals have been made for changing the Federal standard. The current State Standard is .081 ppm. It was established at the same time the original Federal Standard was established and has not been changed. In the past the state standard was interpreted to be .080 ppm but a conversion of the actual 160 ug/m<sup>3</sup> standard to ppm yields .081. Therefore, only hourly averages in excess of .081 ppm are considered exceedances of the state standard.

### 3.4 Monitoring

Ozone was monitored at eleven sites in Maine during 1991 using continuous monitoring equipment of two kinds, either chemiluminescence or ultra-violet absorption analyzers. Maine's ozone monitoring season is limited to April through October due to the weather conditions which are not conducive to ozone formation at other times of the year.

Table 3-1 is the 1991 Data Summary for Ozone. Table 3-2 presents the Ozone Historical Comparisons and Table 3-3 presents the Ozone Trends.

The ozone data has been incorrectly interpreted for the last three years and consequently the trends tables have not reported the ozone data in a consistent format. Starting in 1988 data was stored in the AIRS database instead of the old SAROAD system. Under the SAROAD system reports were generated which indicated the percentiles or frequency distribution of all the hourly data values reported. Reports generated under the AIRS system are now based on percentiles calculated based on each days maximum hourly value. Thus, a 50th percentile of

.050 ppm means that 50% of the days monitored had a maximum hourly value during the day of .050 ppm or less. All of the trend information for ozone in this report have been revised to reflect this method of calculation based on the AIRS report format. If a report format can be developed that will provide percentiles based on total hourly values then that statistic will be used in future reports as it is a better indicator of total exposure to high ozone levels.

TABLE 3 - 1  
 1991 OZONE DATA SUMMARY  
 (Parts Per Million)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>HIGHEST CONCENTRATION</u>	<u>SECOND HIGHEST CONCENTRATION</u>	<u>NUMBER OF VIOLATIONS</u>	
					<u>STATE*</u>	<u>FEDERAL**</u>
CUMBERLAND COUNTY						
Cape Elizabeth	Shelter Site	4815	.145	.141	146	4
HANCOCK COUNTY						
Acadia National Park	McFarland Hill Ranger Station	7138	.128	.125	78	2
Dedham	Bald Mountain	5077	.114	.113	30	0
KENNEBEC COUNTY						
Gardiner	Pray Street School	4789	.132	.123	49	1
KNOX COUNTY						
Isle Au Haut	Isle Au Haut Fire Station	3710	.137	.136	123	5
Port Clyde	Port Clyde Ozone	4541	.137	.135	162	4
PENOBSCOT COUNTY						
Milford	French Settlement Road	4070	.096	.093	6	0
SOMERSET COUNTY						
Skowhegan	Somerset Mill	4828	.116	.112	7	0
WASHINGTON COUNTY						
Jonesport	Public Landing	4373	.120	.117	69	0
YORK COUNTY						
Kennebunkport	Parson's Way	4074	.158	.150	119	4
Isle of Shoals	Appledore Island	1294	.147	.146	40	3

\* Total number of hours minus one greater than .081 ppm.

\*\* Number of days in violation. Not a statistical estimate.

TABLE 3 - 2  
 OZONE HISTORICAL COMPARISONS  
 (1-Hour Concentrations)

CAPE ELIZABETH  
Shelter Site

<u>YEAR</u>	<u>SECOND HIGH</u>	<u># OF STATE VIOLATIONS</u>
1978	.160 PPM	202
1979	.155 PPM	116
1980	.178 PPM	141
1981	.122 PPM	98
1982	.140 PPM	117
1983	.163 PPM	187
1984	.146 PPM	148
1985	.165 PPM	141
1986	.128 PPM	68
1987	.152 PPM	76
1988	.168 PPM	269
1989	.136 PPM	81
1990	.144 PPM	69
1991	.141 PPM	146

KENNEBUNKPORT  
Parson's Way

<u>YEAR</u>	<u>SECOND HIGH</u>	<u># OF STATE VIOLATIONS</u>
1982	.120 PPM	42
1983	.148 PPM	149
1984	.147 PPM	184
1985	.168 PPM	190
1986	.138 PPM	62
1987	.145 PPM	67
1988	.168 PPM	230
1989	.147 PPM	103
1990	.162 PPM	129
1991	.150 PPM	119

DEDHAM  
Bald Mountain

<u>YEAR</u>	<u>SECOND HIGH</u>	<u># OF STATE VIOLATIONS</u>
1985	.123 PPM	72
1986	.106 PPM	34
1987	.117 PPM	35
1988	.185 PPM	241
1989	.105 PPM	41
1990	.119 PPM	68
1991	.113 PPM	30

GARDINER  
Gardiner H. S./Pray Street School

<u>YEAR</u>	<u>SECOND HIGH</u>	<u># OF STATE VIOLATIONS</u>
1980	.117 PPM	54
1981	.122 PPM	31
1982	.120 PPM	56
1983	.140 PPM	99
1984	.112 PPM	89
1985	.133 PPM	84
1986	.110 PPM	17
1987	.112 PPM	25
1988	.145 PPM	142
1989	.118 PPM	47
1990	.107 PPM	35
1991	.123 PPM	49

ACADIA  
McFarland Hill Ranger Station

<u>YEAR</u>	<u>SECOND HIGH</u>	<u># OF STATE VIOLATIONS</u>
1982*	.055 PPM	0
1983	.135 PPM	98
1984	.130 PPM	86
1985	.117 PPM	57
1986	.108 PPM	37
1987	.126 PPM	44
1988	.153 PPM	216
1989	.113 PPM	23
1990	.118 PPM	44
1991	.125 PPM	78

\* Not a complete year.

ISLE AU HAUT  
Isle Au Haut Fire Station

<u>YEAR</u>	<u>SECOND HIGH</u>	<u># OF STATE VIOLATIONS</u>
1986	.107 PPM	26
1987	.151 PPM	87
1988	.127 PPM	111
1989	.115 PPM	35
1990	.131 PPM	55
1991	.136 PPM	123

TABLE 3 - 3  
OZONE TRENDS  
(1-Hour Concentrations)

CAPE ELIZABETH  
Shelter Site

PERCENTILES

<u>Year</u>	<u>10%</u>	<u>50%</u>	<u>90%</u>
1978	.018	.026	.054
1979	.038	.053	.106
1980	.033	.049	.097
1981	.030	.047	.086
1982	.033	.052	.082
1983	.034	.049	.095
1984	.034	.051	.100
1985	.037	.052	.092
1986	.032	.048	.075
1987	.034	.048	.074
1988	.033	.050	.106
1989	.034	.048	.070
1990	.031	.046	.077
1991	.034	.048	.089

KENNEBUNKPORT  
Parson's Way

PERCENTILES

<u>Year</u>	<u>10%</u>	<u>50%</u>	<u>90%</u>
1983	.028	.046	.103
1984	.031	.049	.103
1985*	.038	.056	.098
1986	.033	.048	.077
1987	.033	.046	.074
1988	.035	.052	.119
1989	.036	.052	.085
1990	.035	.050	.089
1991	.038	.050	.088

DEDHAM  
Bald Mountain

PERCENTILES

<u>Year</u>	<u>10%</u>	<u>50%</u>	<u>90%</u>
1985	.030	.050	.083
1986	.028	.043	.068
1987	.030	.044	.065
1988	.030	.047	.083
1989	.031	.045	.070
1990	.030	.046	.071
1991	.025	.043	.070

\* Percentiles calculated using  
70% of the data.

GARDINER  
Gardiner H. S./Pray Street School

PERCENTILES

<u>YEAR</u>	<u>10%</u>	<u>50%</u>	<u>90%</u>
1980	.032	.046	.088
1981	.029	.045	.073
1982	.028	.047	.073
1983	.033	.047	.083
1984	.030	.046	.081
1985	.033	.049	.082
1986	.027	.043	.062
1987	.028	.041	.065
1988	.027	.049	.087
1989	.034	.047	.073
1990	.034	.048	.075
1991*	.031	.044	.074

ACADIA  
McFarland Hill Ranger Station

PERCENTILES

<u>YEAR</u>	<u>10%</u>	<u>50%</u>	<u>90%</u>
1982*	.020	.025	.050
1983	.030	.045	.080
1984	.030	.045	.087
1985	.030	.043	.079
1986	.026	.042	.063
1987	.031	.044	.068
1988	.031	.049	.097
1989	.033	.047	.069
1990	.030	.044	.070
1991	.033	.043	.078

\* Not a complete year.

ISLE AU HAUT  
Isle Au Haut Fire Station

PERCENTILES

<u>YEAR</u>	<u>10%</u>	<u>50%</u>	<u>90%</u>
1986	.024	.040	.077
1987	.033	.045	.078
1988	.028	.058	.122
1989	.025	.036	.069
1990	.028	.048	.076
1991	.033	.048	.088

\* Site relocated to the Pray St. School

## 4. NITROGEN DIOXIDE (NO<sub>2</sub>)

### 4.1 Description and Sources

In its pure state, nitrogen dioxide is a reddish-orange-brown gas with a characteristic pungent odor. It is corrosive and a strong oxidizing agent. Nitrogen dioxide comprises about 10% of the oxides of nitrogen (NOX) that are formed when nitrogen in the air combines with oxygen during high temperature combustion. Most of the rest of the NOX emitted by combustion sources is nitric oxide (NO). However, during the day most of the NO is photochemically transformed into NO<sub>2</sub>. Thus, essentially all the NOX emitted can be assumed to eventually become NO<sub>2</sub>.

### 4.2 Health and Welfare Effects

Exposure to NO<sub>2</sub> affects the delicate structure of lung tissue. High levels cause lung irritation and potential lung damage. Lower levels have been associated with increased respiratory disease. Oxides of nitrogen can cause serious injury to vegetation, including bleaching or death of plant tissue, loss of leaves, and reduced growth rate. NOX also deteriorates fabrics and fades fabric dyes. Nitrate salts formed from nitrogen oxides have been associated with the corrosion of metals. Nitrogen oxides can also reduce visibility.

### 4.3 Standards

The current standard for NO<sub>2</sub> is an annual arithmetic mean (average) value not to exceed .05 ppm. NO<sub>2</sub> is the only gaseous pollutant for which only a long-term (annual average) standard has been established.

### 4.4 Monitoring

Nitrogen Dioxide was monitored at one site in Maine during 1991 using continuous monitoring equipment. Table 4-1 presents the data collected during 1991.

TABLE 4 - 1  
1991 NITROGEN DIOXIDE DATA SUMMARY  
(Parts Per Million)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>ANNUAL ARITHMETIC MEAN</u>
CUMBERLAND COUNTY Portland	Shelter Site	8216	.016

## 5. SULFUR DIOXIDE (SO<sub>2</sub>)

### 5.1 Description and Sources

Sulfur dioxide is a colorless irritating gas having the same pungent odor as a struck match. Most people can detect its taste at a level of about 0.3 to 1 part per million. SO<sub>2</sub> is highly soluble in water, forming sulfurous acid. On a worldwide basis, SO<sub>2</sub> is considered to be one of the major pollution problems. It is emitted mainly from stationary sources that utilize fossil fuels (coal, oil) such as power plants, ore smelters, and refineries.

### 5.2 Health and Welfare Effects

The health effects of sulfur dioxide appear to be always associated with high levels of particulates or other pollutants. The world's major recorded air pollution disasters have been associated with high levels of sulfur dioxide and particulates. The excess deaths attributed to these pollutants were due to respiratory failures and occurred predominantly, but not exclusively, in the elderly and infirm. Atmospheres containing high levels of sulfur dioxide are associated with elevated concentrations of other sulfur compounds such as sulfates and sulfuric acid mists, which are corrosive and potentially carcinogenic.

The corrosiveness of SO<sub>2</sub> and its derivatives also causes crop and material damage. Its transport and transformation into sulfurous and sulfuric acids contribute to acid precipitation, causing soils and lakes to become seriously acidified.

### 5.3 Standards

There are two existing Primary National Ambient Air Quality Standards for sulfur dioxide. The first is a long-term one year arithmetic average of 0.03 parts per million (ppm). The second is a short-term 24-hour average standard where concentrations are not to exceed 0.14 ppm more than once per year. The current Secondary NAAQS for SO<sub>2</sub> is a 3-hour average concentration of 0.5 ppm not to be exceeded more than once per year.

In addition there are three state standards for sulfur dioxide. The first is a long-term one-year arithmetic average of .022 parts per million. The second was a short-term 24-hour average standard of .088 ppm not to be exceeded. The third was a short-term 3-hour average concentration of .439 ppm not to be exceeded. During 1987 both of the short-term standards were amended to allow for one exceedance per year.

### 5.4 Monitoring

Sulfur dioxide was monitored at nineteen sites in Maine during 1991 using continuous monitoring equipment utilizing either the pulsed fluorescent or coulometric methods.



Table 5-1 is the 1991 Data Summary for SO<sub>2</sub>. Tables 5-2 and 5-3 present the SO<sub>2</sub> Historical Comparison Data. Table 5-3 in past years had indicated violations but because one exceedance was allowed per year beginning in 1987 this table now indicates exceedances of the standards rather than violations to maintain continuity for comparisons.

TABLE 5 - 1  
 1991 SULFUR DIOXIDE DATA SUMMARY  
 (Parts Per Million)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>HIGHEST 3-HOUR AVERAGE</u>	<u>SECOND HIGHEST 3-HOUR AVERAGE</u>	<u>HIGHEST 24-HOUR AVERAGE</u>	<u>SECOND HIGHEST 24-HOUR AVERAGE</u>	<u>ANNUAL ARITH. MEAN</u>
ANDROSCOGGIN COUNTY							
Lewiston	Country Kitchen Parking Lot	7851	.064	.057	.026	.024	.006
Livermore Falls	James River/Otis	726	.040	.038	.019	.011	.006
Livermore Falls	Pine State Power	7911	.092	.075	.053	.038	.007
AROOSTOOK COUNTY							
Madawaska	Albert Street	8239	.126	.120	.072	.049	.010
Madawaska	U. S. Post Office	8307	.104	.100	.048	.046	.010
Madawaska	Madawaska High School	8283	.109	.102	.045	.039	.004
CUMBERLAND COUNTY							
Portland	Shelter Site	8261	.058	.058	.039	.033	.009
HANCOCK COUNTY							
Acadia NP	Acadia NP	1870	.019	.017	.013	.008	.002*
Bucksport	240 Main Street	3137	.050	.045	.034	.029	.008*
KENNEBEC COUNTY							
Waterville	Front Street	8255	.109	.076	.042	.035	.009
OXFORD COUNTY							
Mexico	Hunt's Property	8247	.049	.040	.020	.020	.007
Rumford	Taylor Mountain I	8209	.047	.043	.022	.020	.005
Rumford	Taylor Mountain II	8261	.064	.049	.027	.021	.006
Rumford	Village Green Site	8275	.074	.053	.033	.019	.005
PENOBSCOT COUNTY							
Lincoln	Thomas Motel Trailer Park	8022	.101	.085	.042	.035	.005
Millinocket	York Street	8580	.079	.066	.035	.026	.004
East Millinocket	Main Street	5664	.029	.015	.008	.007	.002*
SAGADAHOC COUNTY							
Bath	Coal Pocket Site	5974	.050	.043	.030	.027	.006*
YORK COUNTY							
Biddeford	Biddeford Treatment Plant	5452	.057	.056	.028	.028	.007*

\* Insufficient data collected for valid annual arithmetic mean.

TABLE 5 - 2  
SULFUR DIOXIDE HISTORICAL COMPARISONS  
(Maximum 24-Hour Concentrations of Sulfur Dioxide)

<u>SITE</u>	<u>ADDRESS</u>	<u>MAXIMUM 24-HOUR CONCENTRATION (PPM)</u>						
		<u>1985</u>	<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>	<u>1991</u>
ANDROSCOGGIN COUNTY								
Lewiston	Country Kitchen Parking Lot	.043	.047	.038	.053	.042	.027	.026
AROOSTOOK COUNTY								
Madawaska	Albert Street	.058	.080	.072	.071	.048	.093	.072
Madawaska	U. S. Post Office	.061	.068	.084	.073	.069	.042	.048
Madawaska	Madawaska High School	.037	.046	.076	.057	.032	.027	.045
CUMBERLAND COUNTY								
Portland	Shelter Site	.050	.062	.047	.047	.044	.034	.039
HANCOCK COUNTY								
Acadia National Park	McFarland Hill Ranger Station	--	--	--	--	.011	.007	.013
KENNEBEC COUNTY								
Waterville	Front Street	--	--	--	--	--	.029	.042
OXFORD COUNTY								
Mexico	Hunt's Property	.070	.068	.043	.067	.064	.054	.020
Rumford	Taylor Mountain I	.066	.086	.098	.125	.044	.066	.022
Rumford	Taylor Mountain II	.050	.067	.065	.074	.053	.063	.027
Rumford	Village Green Site	.031	.059	.042	.061	.049	.046	.033
PENOBSCOT COUNTY								
Lincoln	Thomas Motel Trailer Park	.051	.037	.039	.036	.041	.059	.042
Millinocket	York Street	.046	.061	.048	.038	.044	.102	.035
East Millinocket	Main Street	--	--	--	.031	.011	.009	.008
SAGADAHOC COUNTY								
Bath	Coal Pocket Site	--	--	--	--	--	.031	.030
YORK COUNTY								
Biddeford	Biddeford Treatment Plant	--	--	--	.044	.032	.024	.028

\* Not a complete year.

TABLE 5 - 3  
 SULFUR DIOXIDE HISTORICAL COMPARISONS  
 (Sites with exceedances of the standards in the past seven years)

<u>SITES</u>	<u>ADDRESS</u>	<u>TOTAL NUMBER OF EXCEEDANCES*</u>						
		<u>1985</u>	<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>	<u>1991</u>
OXFORD COUNTY								
Rumford	Taylor Mountain I	0	0	1	1	0	0	0
PENOBSCOT COUNTY								
Millinocket	York street	0	0	0	0	0	1	0

\* Includes 3-Hour and 24-Hour Exceedances.

## 6. PARTICULATES (TSP and PM10)

### 6.1 Description and Sources

Particulates is the term given to the tiny particles of solid or semi-solid material found in the atmosphere. It is this "dirt" in the air that is visible as a "Brown Cloud", haze or smog. The sources of particulates are many: wind-blown dust and sand from roadways, fields, and construction; coal dust, fly ash, and carbon black from various combustion sources; and automobile exhaust, to name a few. Particulates that range in size from less than 0.1 micrometer up to approximately 45 micrometers are called "total suspended particulates". Particles larger than that range tend to settle out of the air and not remain suspended, except in high winds.

### 6.2 Health and Welfare Effects

The human nose filters out 99 percent of the large and medium-sized particles. The rest enter the windpipe and lungs, where some, known as inhalable particulates, cling to protective mucous and are removed. Some of the smallest, called respirable particulates, are deposited in the lungs' tiny air sacs (alveoli).

In the lungs particulates slow down the exchange of oxygen with carbon dioxide in the blood, causing shortness of breath. The heart may be strained because it must work harder to compensate for oxygen loss. Usually the people most sensitive to these conditions have respiratory diseases like emphysema, bronchitis, asthma, or heart problems. The elderly and children are also sensitive.

Particles themselves may be poisonous if inhaled or absorbed, damaging remote organs like the kidneys or liver. Swallowed mucous that is laden with poisonous particulate matter may damage the stomach.

In addition, particulates may be carriers of poisonous liquid or gaseous substances. Sulfur dioxide, a major air pollutant in its own right, is frequently absorbed by particulates and can react with them to form sulfates. Sulfates react with moisture in the air or in the respiratory tract to form a corrosive liquid (sulfuric acid) that irritates delicate membranes and slows down the cleansing action of mucous. This effect can reduce the body's ability to remove harmful bacteria, increasing the possibility of infection.

Adverse health effects from particulate matter aren't always seen immediately. Particulates can accumulate in the lungs after repeated, long-term exposure, causing respiratory distress and other health problems that may be manifested later.

Particles in the air block out and scatter sunlight, reducing visibility. Particulates soil and corrode metals, masonry, and textiles. Irritating odors are often associated with particulates, also.

### 6.3 Standards

#### Primary:

At the beginning of 1987 the primary particulate standards were for total suspended particulates (TSP), independent of particle size or chemical composition. The long-term standard was an annual geometric mean not to exceed 75 micrograms of particulates per cubic meter of air (ug/m<sup>3</sup>). The short-term standard was a 24-hour average of 260 ug/m<sup>3</sup> not to be exceeded more than once per year.

In July of 1987 EPA published revised particulate standards to account for the deeper inhalability of small particles and eliminated the total suspended particulate standards. The new standards, rather than applying to TSP, apply to inhalable or fine particulates. A particle size of 10 micrometers was selected as the upper size limit with a 24-hour concentration of 150 ug/m<sup>3</sup> and an annual standard of 50 ug/m<sup>3</sup> expressed as an expected annual arithmetic mean(AAM). The short term standard is attained when the expected number of exceedances is no more than one per year. The expected AAM is determined by averaging the annual arithmetic averages from three successive years of data.

#### Secondary:

The secondary TSP standard was a 24-hour average of 150 ug/m<sup>3</sup> not to be exceeded more than once per year, designed to protect from soiling, corrosion, etc.

When EPA adopted the fine particulate standards they eliminated the secondary TSP standards and made the secondary fine particulate standards equal to the primary fine particulate standards.

#### State Standards:

As of the end of 1988 the State Standards for total suspended particulates still included an annual geometric mean of 60 micrograms per cubic meter and a 24-hour standard of 150 micrograms per cubic meter not to be exceeded. In addition, the Board of Environmental Protection adopted the federal fine particulate standards for both the short term twenty-four hour and the annual arithmetic mean.

In 1989 the State Legislature passed a more restrictive annual standard for fine particulates of 40 ug/m<sup>3</sup>. In addition, the TSP annual state standard was eliminated and the 24 hour standard was changed to be an indicator of a nuisance condition.

### 6.4 Monitoring

Total Suspended Particulates were monitored at 25 sites in Maine during 1991 using High-Volume Particulate Air Samplers (Hi-Vols).

Hi-Vols operate on the same principle as a vacuum cleaner in that the air is drawn through a filter to "catch the dust". The difference is that a Hi-Vol draws a calibrated volume of air through a pre-weighed

filter pad (rather than a bag) for a twenty-four hour period. The change in weight of the filter pad is recorded as total suspended particulate or TSP in micrograms of particulates per cubic meter of air.

Table 6-1 is a summary of the TSP data collected in Maine during 1991. Table 6-2 is a historical comparison of the TSP Annual Geometric Means at sites which have been in existence over the last two years. Table 6-3 summarizes the number of exceedances of the TSP nuisance standard which have occurred over the last six years and the sites at which they occurred.

Fine particulates were monitored at 39 sites during 1991 using PM10 samplers. The sampling is being conducted to obtain data to evaluate the federal and state fine particulate standards, to document compliance with those standards and to obtain background data for new and previously licensed sources. The sampling was conducted with size-selective hi-vols.

The data collected and the sites which were in operation during 1991 have been summarized in Table 6-4. Tables 6-5 and 6-6 provide some historical comparison data over the last few years these monitors have been in operation.

TABLE 6 - 1  
 1991 TOTAL SUSPENDED PARTICULATES DATA SUMMARY  
 (Micrograms Per Cubic Meter)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>HIGHEST 24-HOUR</u>	<u>SECOND HIGHEST</u>	<u>THIRD HIGHEST</u>	<u>ANNUAL GEOMETRIC MEAN</u>
ANDROSCOGGIN COUNTY						
Lewiston	Country Kitchen Parking Lot	58	231	213	210	55.0
CUMBERLAND COUNTY						
Westbrook	Research Building	96	143	125	124	51.0
FRANKLIN COUNTY						
Jay	Crash Road	179	79	63	56	18.0
Jay	Jay Hill	180	82	80	79	21.5
Jay	Burnham	178	145	105	96	28.8
Jay	Weather Level I	178	117	99	94	26.0
KENNEBEC COUNTY						
Waterville	Front Street	57	230	213	162	48.7
Winslow	Gulley Hill Road	162	173	123	119	42.0
Winslow	Boston Avenue	163	147	106	100	32.7
KNOX COUNTY						
Thomaston	Marsh Road	118	111	64	60	21.5
Thomaston	Mitchell Property	120	79	79	73	20.5
OXFORD COUNTY						
Mexico	Labonville's	60	132	114	110	35.9
Rumford	Taylor Mountain I	60	114	78	68	21.7
Rumford	Village Green Site	59	91	76	76	24.3
PENOBSCOT COUNTY						
Bangor	Kenduskeag Pump Station	60	179	168	108	47.8
Brewer	Brewer Junior High School	46	118	108	105	34.7
Lincoln	Thomas Motel Trailer Park	163	252	208	145	37.9
Lincoln	Lincoln Post Office Building	58	157	155	155	41.0
Millinocket	York Street	99	132	100	99	29.5
Old Town	Marsh Island Apartments	47	158	116	103	37.8



TABLE 6 - 1 (continued)  
 1991 TOTAL SUSPENDED PARTICULATES DATA SUMMARY  
 (Micrograms Per Cubic Meter)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>HIGHEST 24-HOUR</u>	<u>SECOND HIGHEST</u>	<u>THIRD HIGHEST</u>	<u>ANNUAL GEOMETRIC MEAN</u>
SOMERSET COUNTY						
Skowhegan	Hinckley	15	39	34	31	13.3
Skowhegan	Eaton Ridge	15	52	40	29	13.1
WASHINGTON COUNTY						
Woodland	Woodland High School	62	116	111	103	26.2*
YORK COUNTY						
Biddeford	Biddeford Treatment Plant	56	108	89	86	32.2
Biddeford	57 Birch Street	102	98	94	86	36.2

\* Insufficient data collected for valid annual geometric mean.

TABLE 6 - 2  
TOTAL SUSPENDED PARTICULATES HISTORICAL COMPARISON  
ANNUAL GEOMETRIC MEANS (UG/M3)

<u>SITE</u>	<u>ADDRESS</u>	<u>ANNUAL GEOMETRIC MEANS (ug/m3)</u>					
		<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>	<u>1991</u>
<b>ANDROSCOGGIN COUNTY</b>							
Lewiston	Country Kitchen Parking Lot	--	--	--	50.5	47.1	55.0
<b>CUMBERLAND COUNTY</b>							
Westbrook	Research Building	67.4	71.2	62.1*	62.0	51.9	51.0
<b>FRANKLIN COUNTY</b>							
Jay	Crash Road	18.9	19.4	20.7	19.6	16.0	18.0
Jay	Jay Hill	24.6	25.1	26.0	25.2	20.9	21.5
Jay	Burnham	--	--	36.0	32.9	28.1	28.8
Jay	Weather Level I	33.5	34.1	38.3	35.0	27.5	26.0
<b>KENNEBEC COUNTY</b>							
Waterville	Front Street Municipal Park	--	--	--	--	39.9	48.7
Winslow	Gulley Hill Road	--	43.6	44.1	51.9*	41.1	42.0
Winslow	Boston Avenue	--	--	--	--	33.3	32.7
<b>KNOX COUNTY</b>							
Thomaston	Marsh Road	23.5	23.4	23.9	23.4	20.5	21.5
Thomaston	Mitchell Property	22.0	21.9	24.5	25.1	21.3	20.5
<b>OXFORD COUNTY</b>							
Mexico	Labonville's	46.6	40.8	43.3	46.5	39.7	35.9
Rumford	Taylor Mountain I	33.0	30.0	30.7	33.8	26.5	21.7
Rumford	Village Green Site	29.7	27.2	27.7	29.7	28.8	24.3
<b>PENOBSCOT COUNTY</b>							
Bangor	Kenduskeag Pump Station	59.4	53.0	56.3	56.2	45.3	47.8
Brewer	Brewer Junior High School	36.5	37.0	37.4	36.8	31.8	34.7
Lincoln	Thomas Motel Trailer Park	34.9	33.9	34.1	33.9	32.8	37.9
Lincoln	Lincoln Post Office Building	34.2	30.3	32.3	35.2	39.0	41.0
Millinocket	York Street	37.3	34.4	33.8	32.1	29.0	29.5
Old Town	Marsh Island Apartments	32.6	36.0	34.6	36.0	30.5	37.8
<b>SOMERSET COUNTY</b>							
Skowhegan	Hinckley	16.6	18.0	14.9	16.8	14.1	13.3

TABLE 6 - 2 (continued)  
 TOTAL SUSPENDED PARTICULATES HISTORICAL COMPARISON  
 ANNUAL GEOMETRIC MEANS (UG/M3)

<u>SITE</u>	<u>ADDRESS</u>	<u>ANNUAL GEOMETRIC MEANS (ug/m<sup>3</sup>)</u>					
		<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>	<u>1991</u>
Skowhegan	Eaton Ridge	17.1	15.5	14.0	18.2	14.0	13.1
WASHINGTON COUNTY							
Woodland	Woodland High School	33.2	29.0	26.5	26.4	34.2*	26.2
YORK COUNTY							
Biddeford	Biddeford Treatment Plant	38.8	36.0	40.8	32.1	27.5	32.2
Biddeford	57 Birch Street	--	--	--	--	34.2	36.2

\* Insufficient data collected for valid annual geometric mean.

TABLE 6 - 3  
 TOTAL SUSPENDED PARTICULATES HISTORICAL COMPARISON  
 (Sites with samples greater than 150 ug/m<sup>3</sup>)

<u>SITE</u>	<u>ADDRESS</u>	<u>TOTAL NUMBER OF SHORT TERM EXCEEDANCES</u>					<u>1991</u>
		<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>	
<b>ANDROSCOGGIN COUNTY</b>							
Lewiston	Country Kitchen Parking Lot	0	0	0	0	0	3
<b>CUMBERLAND COUNTY</b>							
Westbrook	Research Building	15	11	0	4	1	0
<b>FRANKLIN COUNTY</b>							
Jay	Burnham	0	1	1	0	0	0
Jay	Weather Level I	0	1	0	0	0	0
<b>KENNEBEC COUNTY</b>							
Waterville	Front Street	-	-	-	-	3	3
Winslow	Gulley Hill Road	-	2	1	6	2	1
<b>KNOX COUNTY</b>							
Thomaston	Mitchell Property	0	2	3	0	0	0
<b>OXFORD COUNTY</b>							
Mexico	Labonville's	0	0	0	2	0	0
<b>PENOBSCOT COUNTY</b>							
Bangor	Kenduskeag Pump Station	6	2	1	2	0	2
Lincoln	Lincoln Post Office Building	1	2	0	2	3	3
Lincoln	Thomas Motel Trailer Park	0	0	2	4	0	2
Millinocket	York Street	1	4	0	0	0	0
Old Town	Marsh Island Apartments	0	1	1	2	0	1
Woodland	Woodland High School	8	5	0	0	2	0

TABLE 6 - 4  
 1991 FINE PARTICULATE DATA SUMMARY  
 (Micrograms Per Cubic Meter)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>HIGHEST 24-HOUR</u>	<u>SECOND HIGHEST</u>	<u>THIRD HIGHEST</u>	<u>ANNUAL ARITH. MEAN</u>	<u>ANNUAL GEOM. MEAN</u>
ANDROSCOGGIN COUNTY							
Lewiston	Country Kitchen Parking Lot	56	66	66	59	28.5	25.5
AROOSTOOK COUNTY							
Madawaska	Big Daddy's Restaurant	56	95	82	73	32.1	29.1
Presque Isle	Northeastland Hotel	204	123	120	106	27.7	26.7
Presque Isle	Regional Office	115	46	45	36	16.3	14.8
T12R8	Bald Mountain Project	123	35	31	31	14.6	7.5*
CUMBERLAND COUNTY							
Bridgton	Upper Ridge Road	55	44	33	29	12.2	10.7
Portland	Shelter Site	47	54	54	53	24.7	22.6
Portland	Tukey's Bridge	50	85	71	64	27.6	24.9
Westbrook	Research Building	109	59	51	43	22.1	20.4
Westbrook	N. E. T. & T. Company	58	45	39	38	19.1	17.3
FRANKLIN COUNTY							
Jay	Jay Hill	175	57	53	53	19.1	16.3
Jay	Weather Level I	177	45	44	40	16.4	14.6
KENNEBEC COUNTY							
Augusta	Rines Hill Parking Lot	47	69	53	52	26.3	21.7
Waterville	Front Street	58	67	66	62	28.0	24.6
Winslow	Boston Avenue	157	57	52	50	21.6	18.7
KNOX COUNTY							
Thomaston	Marsh Road	115	54	40	35	15.3	13.4
Thomaston	Mitchell Property	119	53	50	42	15.2	12.8
OXFORD COUNTY							
Mexico	Labonville's	178	106	52	51	20.6	17.9
Rumford	Taylor Mountain I	165	105	49	45	17.8	14.8
Rumford	Village Green	178	81	57	43	17.2	14.8
PENOBSCOT COUNTY							
Bangor	Kenduskeag Pump Station	60	51	48	48	25.1	22.9
Brewer	Brewer Junior High School	8	45	29	28	21.4	18.9

TABLE 6 - 4 (continued)  
 1991 FINE PARTICULATE DATA SUMMARY  
 (Micrograms Per Cubic Meter)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>HIGHEST 24-HOUR</u>	<u>SECOND HIGHEST</u>	<u>THIRD HIGHEST</u>	<u>ANNUAL ARITH. MEAN</u>	<u>ANNUAL GEOM. MEAN</u>
Lincoln	Thomas Motel Trailer Park	166	50	49	45	18.2	15.8
Lincoln	Lincoln Street	154	46	37	35	13.1	11.3
Lincoln	Penobscot River	163	64	44	40	11.5	8.9
Lincoln	Lincoln Post Office Building	163	91	78	73	26.8	22.8
Millinocket	Katahdin Nursing Home	40	37	31	29	14.3	12.9
Millinocket	York Street	100	61	60	41	15.5	13.6
East Millinocket	Library/Municipal Building	72	55	38	37	17.5	14.8
Old Town	Marsh Island Apartments	10	40	36	36	21.0	18.1
Orrington	Center Drive School	58	40	34	29	12.8	10.8
Hampden	McGraw School	59	39	34	34	14.4	12.4
SAGADAHOC COUNTY							
Bath	Coal Pocket Site	238	65	58	57	19.7	17.5
SOMERSET COUNTY							
Skowhegan	Hinckley	57	36	34	30	14.2	12.1
Skowhegan	Eaton Ridge	51	38	35	26	14.0	12.4
WASHINGTON COUNTY							
Woodland**	Secondary Treatment Pipeline	57	82	46	43	19.2	16.1
Woodland**	Woodland High School	146	89	84	65	23.3	18.8
Woodland**	Background	59	42	28	28	13.4	11.9
YORK COUNTY							
Biddeford	Biddeford Treatment Plant	57	59	59	47	22.1	20.0

\* Insufficient data collected for valid annual geometric mean.

\*\* Data from this site is under review and may be voided.

TABLE 6 - 5  
 FINE PARTICULATE HISTORICAL COMPARISON  
 ANNUAL ARITHMETIC MEANS (ug/m<sup>3</sup>)

<u>SITE</u>	<u>ADDRESS</u>	<u>ANNUAL ARITHMETIC MEANS (ug/m<sup>3</sup>)</u>					
		<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>	<u>1991</u>
ANDROSCOGGIN COUNTY							
Lewiston	Country Kitchen Parking Lot	--	--	--	--	24.7	28.5
AROOSTOOK COUNTY							
Madawaska	Big Daddy's Restaurant	36.4	31.8	33.4	33.2	34.9	32.1
Presque Isle	Northeastland Hotel	31.0	29.2	26.4	30.0	29.0	27.7
Presque Isle	Regional Office	--	--	--	15.8	14.1	16.3
CUMBERLAND COUNTY							
Bridgton	Upper Ridge Road	16.0	16.2	12.3	11.5	13.4	12.2
Portland	Shelter Site	30.5	30.9	24.4	26.1	22.5	24.7
Westbrook	Research Building	--	--	25.0	24.0	21.5	22.1
Westbrook	N. E. T.&T. Company	--	--	21.0	20.7	17.3	19.1
FRANKLIN COUNTY							
Jay	Jay Hill	21.3	18.9	20.6	22.4	18.2	19.1
Jay	Weather Level I	--	--	17.7	18.1	15.6	16.4
KENNEBEC COUNTY							
Waterville	Front Street	--	--	--	--	25.8	28.0
Winslow	Boston Avenue	--	--	--	--	27.8	21.6
KNOX COUNTY							
Thomaston	Marsh Road	--	--	20.9	17.5	16.3	15.3
Thomaston	Mitchell Property	--	--	22.5	18.2	15.3	15.2
OXFORD COUNTY							
Mexico	Labonvilles	--	30.3	30.5	30.3	24.1	20.6
Rumford	Village Green	--	--	21.1	23.4	19.3	17.2
PENOBSCOT COUNTY							
Bangor	Kenduskeag Pump Station	--	--	30.5	27.3	20.5	25.1
Lincoln	Thomas Motel Trailer Park	30.3	30.8	22.9	23.1	18.9	18.2
Lincoln	Lincoln Street	--	--	--	--	12.7	13.1
Lincoln	Penobscot River	--	--	--	--	11.7	11.5
Lincoln	Lincoln Post Office Building	--	--	--	--	22.5	26.8

TABLE 6 - 5 (continued)  
 FINE PARTICULATE HISTORICAL COMPARISON  
 ANNUAL ARITHMETIC MEANS (ug/m3)

<u>SITE</u>	<u>ADDRESS</u>	<u>ANNUAL ARITHMETIC MEANS (ug/m3)</u>					
		<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>	<u>1991</u>
Millinocket	Katahdin Nursing Home	--	--	--	18.3	15.3	14.3
Millinocket	York street	--	--	16.0	18.9	16.2	15.5
East Millinocket	Library/Municipal Building	--	--	14.4	20.0	16.2	17.5
Orrington	Center Drive school	--	13.9	14.0	13.2	11.5	12.8
Hampden	McGraw School	--	15.3	15.7	15.1	12.9	14.4
<b>SAGADAHOC COUNTY</b>							
Bath	Coal Pocket Site	--	--	--	--	15.9	19.7
<b>SOMERSET COUNTY</b>							
Skowhegan	Hinckley	--	--	22.3	21.9	13.8	14.2
Skowhegan	Eaton Ridge	--	--	14.5	15.5	13.6	14.0
<b>WASHINGTON COUNTY</b>							
Woodland	Secondary Treatment Pipeline	--	--	16.1	17.7	18.5	19.2
Woodland	Woodland High School	21.9	23.4	21.7	21.9	23.7	23.3
Woodland	Background	--	--	10.7	12.7	13.2	13.4
<b>YORK COUNTY</b>							
Biddeford	Biddeford Treatment Plant	--	--	--	26.9	22.0	22.1



TABLE 6 - 6  
 FINE PARTICULATE HISTORICAL COMPARISON  
 (Sites with samples greater than 150 ug/m3)

<u>SITE</u>	<u>ADDRESS</u>	TOTAL NUMBER OF SAMPLES GREATER THAN 150 UG/M3					
		<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>	<u>1991</u>
AROOSTOOK COUNTY							
Madawaska	Big Daddy's Restaurant	0	1	1	0	0	0
Presque Isle	Northeastland Hotel	1	3	0	0	1	0

## 7. LEAD (Pb)

### 7.1 Description and Sources

Lead in the ambient air exists primarily as particulate matter in the inhalable size range. The predominant source of atmospheric lead is from motor vehicles that burn "leaded" gasoline. The lead in gasoline is in the form of tetraethyl lead, an "anti-knock" compound. Other major sources of atmospheric lead are the extraction and processing of metallic ores.

### 7.2 Health and Welfare Effects

When atmospheric lead is breathed in, it is absorbed into the bloodstream and distributed throughout the body along with lead from contaminated food and drinking water. Lead accumulation in the body can impair the production of hemoglobin. Clinical lead poisoning occurs when the body's accumulation of lead becomes too high. Symptoms of lead poisoning range from loss of appetite, fatigue, cramps and constipation, and pains in the ankles and wrists to loss of power in the arms and legs, anemia, kidney disease, mental retardation, blindness and death. Lead concentrations in the ambient air are not sufficient to produce lead poisoning but they do increase the risk of harm when other sources of lead are present. And, indirectly, lead fallout from automotive exhaust onto soil and street surfaces can be ingested in considerable amounts by infants and young children.

### 7.3 Standards

The current National Ambient Air Quality Standard for lead is a 3-month (calendar quarter) average concentration not to exceed 1.5 micrograms of lead per cubic meter of air.

The current State Standard for lead is a 24-hour average concentration of 1.5 micrograms of lead per cubic meter of air not to be exceeded more than once per year.

### 7.4 Monitoring

Lead was monitored at eight sites in Maine during 1991 by taking samples of the Hi-Vol filters from those sites and analyzing the samples for lead content using an atomic absorption analyzer.

Tables 7-1 and 7-2 are the 1991 Data Summaries for Lead. Table 7-3 presents the Lead Historical Comparison Data.

TABLE 7 - 1  
 1991 LEAD DATA SUMMARY  
 (Micrograms Per Cubic Meter)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>HIGHEST 24-HOUR</u>	<u>SECOND HIGHEST</u>	<u>THIRD HIGHEST</u>	<u>ANNUAL GEOMETRIC MEAN</u>
ANDROSCOGGIN COUNTY						
Lewiston	Country Kitchen Parking Lot	58	.05	.05	.03	.01
CUMBERLAND COUNTY						
Portland	Shelter Site	45	.08	.08	.06	.02
Portland	Tukey's Bridge	56	.06	.04	.04	.02
PENOBSCOT COUNTY						
Bangor	Kenduskeag Pump Station	60	.03	.02	.02	.01
YORK COUNTY						
Biddeford	Biddeford Treatment Plant	116	.20	.08	.04	.01
Biddeford	57 Birch Street	104	.05	.04	.02	.01
Saco	Saco Island - CMP	104	.10	.07	.07	.02
Saco	Spring Street	105	.06	.04	.04	.01

TABLE 7 - 2  
 1991 LEAD DATA SUMMARY BY QUARTERS  
 (Micrograms Per Cubic Meter)

<u>SITE</u>	<u>ADDRESS</u>	1991 QUARTERLY AVERAGES			
		<u>1ST</u>	<u>2ND</u>	<u>3RD</u>	<u>4TH</u>
ANDROSCOGGIN COUNTY					
Lewiston	Country Kitchen Parking Lot	.02	.02	.01	.02
CUMBERLAND COUNTY					
Portland	Shelter Site	.03	.02	.02	.03
Portland	Tukey's Bridge	.03	.02	.02	.03
PENOBSCOT COUNTY					
Bangor	Kenduskeag Pump Station	.01	.01	.01	.01
YORK COUNTY					
Biddeford	Biddeford Treatment Plant	.01	.02	.01	.02
Biddeford	57 Birch Street	.02	.01	.01	.02
Saco	Saco Island - CMP	.02	.03	.01	.02
Saco	Spring Street	.02	.02	.01	.02

TABLE 7 - 3  
 LEAD HISTORICAL COMPARISONS  
 (Micrograms Per Cubic meter)

<u>SITE</u>	<u>ADDRESS</u>	<u>MAXIMUM 24-HOUR CONCENTRATION / AAM</u>						
		<u>1985</u>	<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>	<u>1991</u>
<b>ANDROSCOGGIN COUNTY</b>								
Lewiston	Country Kitchen Parking Lot	-----	-----	-----	-----	0.12/0.03	0.04/0.02	0.05/0.02
<b>CUMBERLAND COUNTY</b>								
Portland	Shelter Site	0.53/0.19	0.33/0.11	0.27/0.07	0.17/0.06	0.10/0.04	0.12/0.03	0.08/0.03
Portland	Tukey's Bridge	1.10/0.42	0.87/0.35	-----	-----	0.08/0.04	0.08/0.03	0.06/0.02
<b>PENOBSCOT COUNTY</b>								
Bangor	Kenduskeag Pump Station	0.64/0.15	0.18/0.07	0.12/0.04	0.08/0.03	0.09/0.02	0.10/0.02	0.03/0.01
<b>YORK COUNTY</b>								
Biddeford	Biddeford Treatment Plant	-----	-----	-----	-----	0.82/0.06	0.63/0.03	0.20/0.01
Biddeford	57 Birch Street	-----	-----	-----	-----	-----	0.26/0.02	0.05/0.01
Saco	Saco Island	-----	-----	-----	-----	-----	0.70/0.03	0.10/0.02
Saco	Spring Street	-----	-----	-----	-----	-----	0.07/0.02	0.06/0.02

## 8. SULFATES (SO<sub>4</sub>) AND NITRATES (NO<sub>3</sub>)

### 8.1 Description and Sources

Sulfates and Nitrates are compounds of varying harmfulness found everywhere in the atmosphere. They are produced by nature as well as man. Man-made sulfates have their origin in sulfur dioxide while nitrates have theirs in nitrogen oxides. Fine particulate compounds, including sulfates and nitrates are formed from chemical reactions between sulfur dioxide or nitrogen dioxide emitted into the air and other substances present there. These fine particulate compounds have a long atmospheric residence time, can be transported in the air for long distances, and are capable of penetrating deeply into the human respiratory tract.

### 8.2 Health and Welfare Effects

Epidemiological studies of populations exposed to particulate sulfates have shown that atmospheric sulfates, more than sulfur dioxide gas or total suspended particulates, are related to aggravation of asthma, aggravation of heart and lung disease in the elderly, and impairment of lung function in school children. This evidence was obtained from EPA's Community Health and Environmental Surveillance System (CHESS). From these studies, estimates of the sulfate threshold for adverse health effects have been derived, as shown in Table 8-1. However, these epidemiological studies have not been substantiated by laboratory studies.

Both sulfates and nitrates are considered to be contributors to the acid deposition problem.

### 8.3 Standards

There are currently no standards for levels of sulfates in ambient air. EPA has been working on a standard and is expected to make a proposal in the future.

There are no standards for nitrates nor are there any proposed.

### 8.4 Monitoring

Sulfate levels were measured at four sites in Maine during 1991 by taking samples of the Hi-Vol filters from those sites and analyzing the samples for sulfates using the Automated Technicon II Methylthymol Blue Procedure. There is no standard yet and the monitoring methodology is questionable but the data is being included in this report as an aid to those interested in further information about Maine's air quality. Table 8 -2 summarizes the sulfate data collected during 1991.

Nitrate levels were not measured in Maine during 1991.

TABLE 8-1

SULFATE THRESHOLDS FOR ADVERSE HEALTH EFFECTS

<u>ADVERSE HEALTH EFFECT</u>	<u>THRESHOLD CONCENTRATION FOR SUSPENDED SULFATES</u>
Aggravation of Asthma	6 to 10 Micrograms Per Cubic Meter for 24 Hours.
Aggravation of Heart and Lung Disease in the Elderly	9 Micrograms Per Cubic Meter for 24 Hours
Subtle Decreases in Childhood Lung Function	9 to 13 Micrograms Per Cubic Meter for 1 Year.
Increase in Acute Respiratory Disease in Children	13 Micrograms Per Cubic Meter for 1 Year.

TABLE 8 - 2  
 1991 SULFATE DATA SUMMARY  
 (Micrograms Per Cubic Meter)

<u>SITE</u>	<u>ADDRESS</u>	<u>NUMBER OF OBSERVATIONS</u>	<u>HIGHEST 24-HOUR</u>	<u>SECOND HIGHEST</u>	<u>THIRD HIGHEST</u>	<u>ANNUAL ARITHMETRIC MEAN</u>
CUMBERLAND COUNTY						
Bridgton	Upper Ridge Road	55	21.3	15.6	12.0	3.6
Portland	Shelter Site	47	22.1	11.1	10.7	3.6
South Portland	SMVTI	28	10.6	10.4	7.3	4.0
OXFORD COUNTY						
Rumford	Taylor Mountain I	60	22.8	20.5	13.7	4.9



## **9. ATMOSPHERIC DEPOSITION**

### **9.1 Description and Sources**

As a result of the combustion of tremendous quantities of fossil fuels such as coal and oil, the United States annually discharges approximately 50 million metric tons of sulfur and nitrogen oxides into the atmosphere. Through a series of complex chemical reactions these pollutants can be converted into acids, which may return to earth as components of either rain or snow. This atmospheric deposition, more commonly known as acid rain, may have severe ecological impacts on widespread areas of the environment.

### **9.2 Health and Welfare Effects**

While direct health effects from acid rain have not been documented there are numerous indirect effects which could have definite effect on mankind. Atmospheric deposition is known to leach heavy metals such as mercury from rocks causing possible contamination of water supplies. Hundreds of lakes in North America and Scandanavia have become so acidic that they can no longer support fishlife. The rain falling on forests and other non-farmlands could, in time, cause extensive changes in the soil chemistry. There is not enough information yet to make it possible to say exactly what the results might be, but there is no reason to think the changes will be beneficial.

### **9.3 Standards**

There are no standards in effect or proposed for atmospheric deposition. The only permanent solution to the acid rain problem is to keep the acid levels low. The only practical way of achieving this is by reducing emissions at their sources.

### **9.4 Monitoring**

During 1991 there were four sites collecting data on atmospheric deposition. Those four sites included two Bureau maintained sites in Bridgton and Acadia National Park, a University of Maine maintained site in Greenville and a National Weather Service maintained site in Caribou. The samples from these four sites are normally collected every Tuesday morning at 9:00 a.m.. Consequently, the samples are not necessarily a single storm event but are more likely to be a composite of all storm events during the previous week. The samples, if there was a significant storm, are used for field measurements of pH and conductivity and are then packaged up for shipment to the National Atmospheric Deposition Program central laboratory in Illinois. In the central laboratory they are also tested for pH and conductivity as well as additional components. Table 9-1 is a summary of the measurements taken at the central laboratory in Illinois from the DEP operated and/or maintained sites and the University of Maine operated site in Greenville for the year 1991. The sulfate deposition figures were corrected for marine aerosol contribution.

TABLE 9 - 1  
1990 ATMOSPHERIC DEPOSITION DATA SUMMARY

<u>SITE</u>	<u>ADDRESS</u>	<u>MAXIMUM*</u>	<u>pH</u>		<u>MEAN**</u>	<u>DEPOSITION (Kg/ha)</u>	
			<u>MINIMUM*</u>			<u>SO4***</u>	<u>NO3</u>
DOWNEAST AIR QUALITY CONTROL REGION (109)							
Acadia National Park	McFarland Hill Ranger Station	5.5	4.0		4.6	14.0	10.0
METROPOLITAN PORTLAND AIR QUALITY CONTROL REGION (110)							
Bridgton	Upper Ridge Road	5.7	3.9		4.6	13.0	9.0
NORTHWEST MAINE AIR QUALITY CONTROL REGION (111)							
Greenville	Squaw Brook	5.6	4.0		4.7	10.0	9.0

\* Lab measurements.

\*\* Precipitation weighted mean.

\*\*\* Corrected for marine aerosol and normalized to 52 weeks.

## 10. HYDROCARBONS (HC)

### 10.1 Description and Sources

Hydrocarbons are a class of compounds containing carbon and hydrogen in various combinations. They are found especially in petroleum, natural gas and coal. Some are gaseous, some liquid and some are solid. There are in fact over a thousand hydrocarbon compounds. Many of the polluting hydrocarbons are discharged into the air by incomplete combustion of organic materials. A major source of this kind of hydrocarbon emission is the burning of gasoline in automobiles. Other major contributors are organic solvent evaporation, industrial processes, solid waste disposal and fuel combustion in stationary sources. The control of hydrocarbon emissions are accomplished by combustion process optimization, recovery by mass transfer principles, restriction of evaporative loss and process material and fuel substitution.

### 10.2 Health and Welfare Effects

Hydrocarbon air pollutants enter into and promote the formation of photochemical smog (ozone) and thus contribute to the development of eye irritation and respiratory tract problems. By themselves, hydrocarbons may induce adverse health effects, although there is relatively little quantitative data to relate individual hydrocarbons to the risk of human disease.

### 10.3 Standards

The present State Standard for non-methane hydrocarbons is a three hour average concentration of 160 ug/m<sup>3</sup>.

### 10.4 Monitoring

Hydrocarbons were not monitored as part of the state's continuous air monitoring program during 1991.